


Enhancing Coordination of Behavioral Health Services after Superstorm Sandy: Planning for Future Disasters

Identifying at-Risk Communities



Initial Data Profile: Atlantic County

Demographics, Behavioral Health Conditions, and Utilization of Health Services (Medicare Fee-for-Service Beneficiaries)

January 7, 2014

This page left intentionally blank

TABLE OF CONTENTS

| | |
|---|----|
| Preface | 1 |
| Introduction | 2 |
| HQSI Project Team | 5 |
| Acknowledgements | 6 |
| Executive Summary | 7 |
| Demographics | 9 |
| • Total Medicare FFS Beneficiary Population by County | 9 |
| • Percent of Medicare FFS Beneficiary Population by Gender by County..... | 10 |
| • Percent of Medicare FFS Beneficiary Population by Race by County..... | 11 |
| • Percent of Medicare FFS Beneficiary Population by Age by County | 12 |
| • Socioeconomic Status by County..... | 12 |
| Behavioral Health Conditions | 13 |
| • Prevalence and Incidence..... | 13 |
| – Summary | 14 |
| – Depression or Proxy Disorders | 15 |
| – Depression..... | 20 |
| – Anxiety Disorders | 23 |
| – Adjustment Disorders | 23 |
| – Post-Traumatic Stress Disorder..... | 24 |
| – Alcohol or Substance Abuse | 27 |
| – Substance Abuse..... | 30 |
| – Suicide and Intentional Self-Inflicted Injury..... | 33 |
| • Risk Factors for Depression or Proxy Disorders..... | 36 |
| – Top Five Risk Factors for Depression or Proxy Disorders..... | 36 |
| – Alzheimer's Disease and Related Disorders or Senile Dementia..... | 39 |
| – Sleep Disturbance..... | 39 |
| – Substance or Alcohol Abuse or Tobacco Use | 40 |
| – Hip/Pelvic Fractures | 40 |
| – Amputations | 41 |
| Utilization | 43 |
| • Outpatient Behavioral Health Services | 43 |
| – Summary | 43 |
| – Assessments | 45 |
| › Depression Screening | 45 |

TABLE OF CONTENTS

| | |
|--|-----------|
| › Diagnostic Psychological Tests | 49 |
| › Health and Behavior Assessment/Intervention..... | 50 |
| › Neuropsychological Tests | 51 |
| › Psychiatric Diagnostic Procedures..... | 53 |
| – Therapies | 55 |
| › Individual Psychotherapy..... | 55 |
| › Family Psychotherapy..... | 57 |
| › Group Psychotherapy..... | 58 |
| › Electroconvulsive Therapy | 59 |
| › Biofeedback Therapy..... | 60 |
| • Inpatient Services | 61 |
| – Summary | 61 |
| – Psychiatric Hospital Admissions..... | 63 |
| – Acute Care Hospitals | 64 |
| › Admissions | 64 |
| › Observation Stays..... | 65 |
| › Emergency Department Visits..... | 66 |
| › 30-Day Hospital Readmissions..... | 67 |
| › Observation Stays Within 30 Days of Discharge | 68 |
| › Emergency Department Visits Within 30 Days of Discharge | 69 |
| – Other Settings..... | 70 |
| › Summary | 70 |
| › Home Health Agency Services | 71 |
| › Skilled Nursing Facility Services | 72 |
| › Hospice Services | 73 |
| › Medical Rehabilitation Services | 74 |
| Appendices | 75 |
| • Appendix A: Behavioral Health Conditions..... | 75 |
| • Appendix B: Risk Factors for Depression or Proxy Disorders..... | 77 |
| • Appendix C: Utilization of Outpatient Mental Health Services | 81 |
| • Appendix D: Utilization of Services – Inpatient and Other Settings | 83 |
| • Appendix E: Time Frames and Formulae | 85 |
| • Appendix F: References..... | 86 |
| Index of Figures | 87 |

PREFACE

On October 29, 2012, Superstorm Sandy hit the Eastern Seaboard, impacting more than a dozen states. New Jersey, which took the brunt of the storm along its densely populated coastline, was devastated. Thousands of residents were displaced, their homes and communities damaged or destroyed.

Lessons learned from prior natural disasters showed that victims of storms like Superstorm Sandy are often at an elevated risk for acute or chronic behavioral health issues such as post-traumatic stress disorder (PTSD), depression, suicide, and alcohol abuse. While disaster-related issues subside over time, evidence shows that victims can experience a prolonged period of elevated risk, especially those with pre-existing chronic mental health issues. Older adults and disabled residents with chronic mental health conditions are at increased risk of deteriorating health, depression, increased isolation, and breakdown in the continuum of health care. Additionally, past natural disasters also show that access to informational resources on disaster-related mental health disorders, outcomes, and service utilization are important factors to consider.

This initial county profile – one of 10 being created for each of the Federal Emergency Management Agency (FEMA)-declared disaster counties in New Jersey – explores potential county and community level health status and health determinants of post-disaster spikes in behavioral health issues and treatments. Subsequent county profiles, to be produced in spring 2014, will include more comprehensive post-Sandy data and an analytic treatment of the predictive value of the initial county profiles in planning for and coordinating post-disaster response resources.

INTRODUCTION

Enhancing Coordination of Behavioral Health Services after Superstorm Sandy: Planning for Future Disasters is a Special Innovation Project funded by the Centers for Medicare & Medicaid Services (CMS). As part of this project, Healthcare Quality Strategies, Inc. (HQSI), the CMS quality improvement organization (QIO) for New Jersey, studied data on prevalence and incidence of selected behavioral health conditions, the utilization of health services, and demographic information from the Medicare claims database for Medicare Fee-for-Service (FFS) beneficiaries residing in the 10 New Jersey FEMA-declared disaster counties after Superstorm Sandy. These counties include Atlantic, Bergen, Cape May, Essex, Hudson, Ocean, Middlesex, Monmouth, Somerset, and Union.

From its analysis, HQSI created data profiles for each of these FEMA-designated counties. This is the initial profile for Atlantic County. Separate county profiles have been developed for each of the FEMA-declared disaster counties. The county profiles can be used to determine and compare the baseline prevalence of the selected behavioral health conditions and utilization of services among all 10 FEMA-declared disaster counties pre-Superstorm Sandy.

In January 2014, additional profiles will be released that will feature a subset of 10 communities. These communities were selected because they had high rates of Medicare FFS beneficiaries both with and at risk for depression or proxy disorders. The community profiles can be used to determine and compare the baseline prevalence of the selected behavioral health conditions and utilization of services in the selected communities compared to their counties.

The county and community profiles are based on Medicare FFS claims data and provide a glimpse into the prevalence and incidence of selected behavioral health conditions and risk factors for depression, as well as the utilization of Medicare-covered behavioral health services among Medicare beneficiaries residing in the selected counties or communities before and after Superstorm Sandy. Since patients with behavioral health conditions may receive other health services because of medical problems caused by their behavioral health conditions or they may avoid utilizing behavioral health services, this report also looks at the utilization of non-behavioral health services.

These profiles are being shared with state and local governments and agencies, healthcare providers, community-based organizations, and the research community to support a community-based approach to enhance the coordination of behavioral health services after a natural disaster, and to increase utilization of the Medicare depression screening benefit which became a covered service in January 2012.

METHODOLOGY

Each county profile compares one county's statistics to the aggregate of the 10 counties and to the other nine counties. Primary data sources include Medicare FFS Part A and Part B claims, the Medicare enrollment database and U.S. Census data. The Medicare enrollment database includes basic demographic statistics such as age, gender, and race while the U.S. Census data provides a proxy indicator (average household income) for socio-economic status. Based on the ICD-9-CM (International Classification of Disease, Ninth Revision, Clinical Modification), CPT (Current Procedural Terminology) or HCPCS (Healthcare Common Procedure Coding System) codes in Medicare Part A and Part B claims, beneficiaries were identified for chronic conditions including diseases/conditions related to behavioral health, such as depression. Appendices A through F contain documentation, technical notes, codes, algorithms, data sources, and references.

Medicare Part A claims were also used to analyze utilization of health services in or by acute care hospitals, skilled nursing facilities, medical rehabilitation facilities, home health agencies, hospice, and inpatient psychiatric facilities. Medicare Part A and Part B claims provide information on the utilization of mental health outpatient services for assessment (e.g., depression screening, diagnostic psychological tests) and treatment (e.g., individual psychotherapy, biofeedback therapy).

To identify beneficiaries with an elevated risk of depression after the storm, HQSI conducted a literature review on risk factors for depression (see Appendix B). Previous studies identified psychosocial and biological factors, increased age, history of cancer, Parkinson's disease, Alzheimer's disease, changes in mental function, and medication side effects as risk factors for developing depression. Based on findings from the literature review and factors available through Medicare claims, logistic regression analysis was conducted with Medicare claims, and the top five risk factors (Alzheimer's disease and related disorders or senile dementia, hip/pelvic fractures, amputations, substance or alcohol abuse or tobacco use, and sleep disturbance) were used to identify beneficiaries with high risk for developing depression or proxy disorders.

MEASUREMENT TIME FRAMES

This profile includes data from January 1, 2011 through March 31, 2013. October 1, 2012 through December 31, 2012 (Q4 2012) is defined as the quarter during which Superstorm Sandy occurred. The post-storm quarter is defined as Q1 2013 (January 1 – March 31, 2013). Results are presented using three different measurement time frames as follows:

- The pre-Sandy time period was defined as January 1, 2011 through September 30, 2012. Statistics on demographics, prevalence of behavioral health conditions and utilization of health services are presented for this 21-month period. These statistics allowed for comparison across affected counties prior to Superstorm Sandy.

- Annual prevalence with rolling quarters of behavioral health conditions and utilization statistics are included to adjust for seasonal variation and to examine possible changes pre- and post-Superstorm Sandy. The time period used for this analysis was January 1, 2011 through March 31, 2013. This time period includes six data time points.
- Quarterly new incidence of the behavioral health conditions that includes five quarters of data from Q1 2012 (January 1, 2012 – March 31, 2012) through Q1 2013 (January 1, 2013 – March 31, 2013) allows the identification of new cases in a quarter when compared to the prior year. It also allows identification of possible changes after the storm when comparing Q1 2013 data against Q1 2012.

DATA CONSIDERATIONS

The available data relating to behavioral health issues as a result of Superstorm Sandy are new, given that the disaster occurred recently. Currently, there is only one quarter of post-storm data available. To examine possible changes, profiles will be updated in 2014 (when another quarter of post-storm data will be available). Claims data processing lag (at least six months), coupled with the one-year project time frame, reduces the optimal time frame for more accurate estimation of post-Sandy effects.

Identification of beneficiaries with behavioral health conditions is based on diagnoses being reported in Medicare FFS claims and could result in underestimation. There is currently no accurate way to identify when certain health conditions began and ended.

According to the subject matter experts consulted for this project, unlike other conditions, behavioral health issues are often underdiagnosed in our society and the stigma associated with behavioral health conditions may prevent people from seeking care in mental health facilities.

This type of county profile can be used to provide a baseline for the prevalence and incidence rates of eight selected behavioral health conditions (see page 13) based on the ICD-9-CM codes through the Medicare claims database. Possibly, after further data collection and analytic development using post-Sandy data, it can also be used to prioritize and plan community and county preparation for the care, tracking, and monitoring of Medicare beneficiary behavioral health status and health care utilization patterns.

HQSI will produce updated profiles in spring 2014 that will include additional data for the post-Superstorm Sandy time period.

Martin P. Margolies
Chief Executive Officer

Mary Jane Brubaker, MCIS
Chief Operating Officer

Diane Babuin, MS, CPHQ
Director, Quality Improvement and Communications

Ya-ping Su, PhD
Director, Research and Analysis

Suzanne Dalton, RN, BS, EdM
Project Manager

Andrew Miller, MD, MPH
Medical Director

Mona Abdalla, BA
Administrative Associate

Zhengyu Bu, MS
Health Services Research Analyst

Sue Chen, MS
Health Services Research Analyst

Wei-Yi Chung, MS
Health Services Research Analyst

Dawn Cullen, BA
Communications Specialist

Kim Karnell, BS
Information Specialist

Janet Knoth, BS, RN, CHPN
Quality Improvement Specialist

Sai Loganathan, PhD
Health Services Research Analyst

Judy Miller, MS, RN
Quality Improvement Specialist

Barbara Perzyna, BS
Visual Communications Specialist

Ziphora Sam, MPH
Health Services Research Analyst

Marianne Sagarese, BSN, RN
Quality Improvement Specialist

Nicole Skyer-Brandwene, MS, R.Ph, BCPS
Quality Improvement Specialist

Ashley Strain, BA
Communications Specialist

ACKNOWLEDGEMENTS

Special thanks to the Subject Matter Experts who assisted with the project by providing feedback and guidance to the HQSI project team.

Carol Benevy, MSW

New Jersey Hope and Healing Project
Barnabas Health Institute for Prevention

Mary Ditri, MA, CHCC

New Jersey Hospital Association

Adrienne Fessler-Belli, MSW, LCSW

New Jersey Department of Human Services
Disaster & Terrorism Branch

Mark Firth, MA, MSW

New Jersey Department of Human Services
Division of Mental Health and Addiction Services

Mary Goepfert, MPA, APR, CPM

New Jersey Group for Access and Integration Needs in Emergencies and Disasters

Connie Greene, MA, CAS, CSW, CPS

Barnabas Health Institute for Prevention

Bob Kley

Mental Health Association in New Jersey, Inc.

Lynn Kovitch, MEd

New Jersey Department of Human Services
Division of Mental Health and Addiction Services

Karen McCoy, RN, BSN

Home Care Association of New Jersey

Elyse Perweiler, MPP, RN

NJ Institute for Successful Aging

Lynn Stefanowicz, MA, LCSW

Meridian Behavioral Health

Megan Sullivan, LPC, LCADC, DRCC

New Jersey Department of Human Services
Disaster & Terrorism Branch

Pete Summers

The New Jersey Association of County and City Health Officials (NJACCHO)

Vito Veneruso

New Jersey Primary Care Association


KEY OBSERVATIONS

1. In the 21 months prior to Superstorm Sandy, 27.0% of Medicare FFS beneficiaries (269.50 per 1,000 beneficiaries) in Atlantic County experienced depression or proxy disorders (anxiety and adjustment disorders).
2. In the 21 months prior to Superstorm Sandy, 18.6% of Medicare FFS beneficiaries (185.94 per 1,000 beneficiaries) in Atlantic County were at risk for depression or proxy disorders.
3. In the 21 months prior to Superstorm Sandy, Atlantic County had the highest prevalence rate of alcohol or substance abuse (60.37 per 1,000 Medicare FFS beneficiaries), substance abuse (34.67 per 1,000 beneficiaries), and suicide and intentional self-inflicted injury (9.24 per 1,000 beneficiaries) among all 10 counties.
4. In the 21 months prior to Superstorm Sandy, Atlantic County had the lowest prevalence rate of Alzheimer's disease and related disorders or senile dementia among all 10 counties.
5. Atlantic County had the largest decrease in the yearly prevalence of PTSD (8.1%) and alcohol or substance abuse (9.2%).
6. The utilization rate of the Medicare depression screening benefit was low among all 10 counties. In Atlantic County the rate was 1.12 per 1,000 Medicare FFS beneficiaries.
7. The most frequently used behavioral health assessment service in Atlantic County was psychiatric diagnostic procedures (9.1%).
8. The most frequently used behavioral health therapy in Atlantic County was individual psychotherapy (6.9%).
9. Atlantic County had the highest utilization rate of group psychotherapy (0.78%) among all 10 counties.
10. Atlantic County had the highest rate of acute care hospital admissions (642.49 per 1,000 Medicare FFS beneficiaries), emergency department visits (510.52 per 1,000 beneficiaries), and emergency department visits within 30 days of discharge (167.45 per 1,000 beneficiaries).
11. Atlantic County had the lowest utilization rate of skilled nursing facility services (9.9%) among all 10 counties.

The *Snapshot of Atlantic County* (Figure 1) summarizes the prevalence of the behavioral health conditions, as well as risk factors for depression or proxy disorders, analyzed for this county profile. This *Snapshot* also lists the most frequently performed behavioral health assessments and therapies in Atlantic County compared to the average among all 10 counties included in this project. The non-behavioral health utilization measures that were calculated for this profile are not included in the *Snapshot*.


| Figure 1. Snapshot of Atlantic County | | |
|--|--|--------------------------|
| Behavioral Health Disorders | Prevalence per 1,000 Beneficiaries (21 Months Prior to Superstorm Sandy) | |
| | Atlantic County | 10 County Average |
| Depression or Proxy Disorders | 269.50 | 251.86 |
| • Depression alone | 171.36 | 164.75 |
| • Anxiety Disorders alone | 163.83 | 144.53 |
| • Adjustment Disorders alone | 45.32 | 43.45 |
| PTSD | 7.13 | 5.36 |
| Alcohol or Substance Abuse | 60.37 | 43.26 |
| • Substance Abuse alone | 34.67 | 23.75 |
| Suicide and Intentional Self-Inflicted Injury | 9.24 | 6.80 |
| Top Five Risk Factors for Depression or Proxy Disorders* | 185.94 | 175.56 |
| • Alzheimer's Disease and related disorders or Senile Dementia | 42.31 | 49.25 |
| • Sleep Disturbance | 35.09 | 32.26 |
| • Substance or Alcohol Abuse or Tobacco Use | 119.73 | 104.94 |
| • Hip/Pelvic Fractures | 9.43 | 10.99 |
| • Amputations | 1.55 | 1.50 |
| Behavioral Health Services | Utilization per 1,000 Beneficiaries (21 Months Prior to Superstorm Sandy) | |
| | Atlantic County | 10 County Average |
| Assessments | | |
| • Depression Screening | 1.12 | 4.81 |
| • Psychiatric Diagnostic Procedures | 90.99 | 81.61 |
| • Neuropsychological Test | 12.04 | 14.18 |
| Therapy | | |
| • Individual Psychotherapy | 69.00 | 69.31 |
| • Family Psychotherapy | 4.02 | 5.09 |
| • Group Psychotherapy | 7.83 | 3.88 |
| Psychiatric Hospital Admissions | 10.58 | 12.56 |

* The top five risk factors were identified based on findings from a literature review (Appendix B) and factors available through Medicare claims. Logistic regression analysis was conducted with Medicare claims.

| Medicare FFS Demographics | Atlantic County | |
|---|-----------------|--|
| <p>At A Glance</p> <p>Total Medicare FFS Population – 51,469</p> <hr/> <p>Females – 28,294 (54.97%)</p> <p>Males – 23,175 (45.03%)</p> <hr/> <p>White (78.45%)</p> <p>Black (14.09%)</p> <p>Asian (2.90%)</p> <p>Hispanic (2.50%)</p> <p>Other (2.06%)</p> <hr/> <p>Average Age 69.06</p> <hr/> <p><i>Source: Medicare Claims Database</i></p> | |  |

TOTAL MEDICARE FFS BENEFICIARY POPULATION BY COUNTY

The total Medicare FFS beneficiary population of Atlantic County is 51,469 (Figure 2).



PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY GENDER BY COUNTY

Females make up 54.97% of the entire Medicare FFS population in Atlantic County and males 45.03% (Figure 3).

| Figure 3. Percent of Medicare FFS Beneficiary Population by Gender by County (January 1, 2011 – March 31, 2013) | | |
|---|------------------|--------------------|
| County | Percent of Males | Percent of Females |
| Atlantic | 45.03 | 54.97 |
| Bergen | 43.14 | 56.86 |
| Cape May | 46.10 | 53.90 |
| Essex | 42.89 | 57.11 |
| Hudson | 42.80 | 57.20 |
| Middlesex | 43.98 | 56.02 |
| Monmouth | 43.89 | 56.11 |
| Ocean | 43.01 | 56.99 |
| Somerset | 43.45 | 56.55 |
| Union | 43.04 | 56.96 |
| Average of 10 counties* | 43.47 | 56.53 |

*Computing the average of all 10 counties in this table will not equal the average shown, as some beneficiaries moved from one county to another during this time frame.

PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY RACE BY COUNTY

The majority of this population is White (78.45%) followed by Black (14.09%), Asian (2.90%), Hispanic (2.50%), and Other (2.06%) (Figure 4).

| Figure 4. Percent of Medicare FFS Beneficiary Population by Race by County (January 1, 2011 – March 31, 2013) | | | | | |
|---|-------------------|-------------------|----------------------|-------------------|------------------|
| County | Percent of Whites | Percent of Blacks | Percent of Hispanics | Percent of Asians | Percent of Other |
| Atlantic | 78.45 | 14.09 | 2.50 | 2.90 | 2.06 |
| Bergen | 83.43 | 4.85 | 2.46 | 4.57 | 4.69 |
| Cape May | 94.48 | 3.63 | 0.40 | 0.40 | 1.09 |
| Essex | 54.00 | 36.02 | 4.24 | 1.81 | 3.93 |
| Hudson | 61.15 | 12.20 | 14.82 | 5.29 | 6.54 |
| Middlesex | 77.29 | 7.65 | 3.48 | 6.26 | 5.32 |
| Monmouth | 87.67 | 7.01 | 0.80 | 1.55 | 2.97 |
| Ocean | 95.47 | 1.98 | 0.56 | 0.56 | 1.42 |
| Somerset | 82.59 | 6.77 | 1.37 | 4.36 | 4.91 |
| Union | 69.39 | 19.68 | 5.52 | 1.84 | 3.58 |
| 10 counties* | 78.42 | 11.21 | 3.49 | 3.08 | 3.80 |

*Computing the average of all 10 counties in this table will not equal the average shown, as some beneficiaries moved from one county to another during this time frame.

PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY AGE BY COUNTY


The beneficiary population in Atlantic County varies by age group with the largest group between ages 65 and 74 years old followed by beneficiaries below 65 years old. The average age of beneficiaries in this county is 69.06, the youngest average age among all 10 counties (Figure 5).

| Figure 5. Percent of Medicare FFS Beneficiary Population by Age by County (January 1, 2011 – March 31, 2013) | | | | | Average Age by County (January 1, 2011 – March 31, 2013) |
|--|--------------|--------------|--------------|--------------|--|
| County | <65 | 65 – 74 | 75 – 84 | 85 and Above | |
| Atlantic | 29.99 | 38.06 | 22.08 | 9.87 | 69.06 |
| Bergen | 21.57 | 38.27 | 26.67 | 13.49 | 72.08 |
| Cape May | 26.28 | 39.31 | 24.02 | 10.40 | 70.20 |
| Essex | 29.31 | 36.96 | 22.34 | 11.39 | 69.46 |
| Hudson | 28.69 | 37.76 | 23.29 | 10.26 | 69.55 |
| Middlesex | 26.36 | 37.27 | 24.57 | 11.80 | 70.51 |
| Monmouth | 26.33 | 38.08 | 23.62 | 11.97 | 70.59 |
| Ocean | 22.85 | 37.89 | 26.50 | 12.75 | 71.51 |
| Somerset | 23.73 | 39.98 | 24.28 | 12.01 | 71.15 |
| Union | 26.28 | 36.41 | 24.23 | 13.09 | 70.77 |
| 10 counties* | 25.56 | 37.86 | 24.56 | 12.02 | 70.72 |

*Computing the average of all 10 counties in this table will not equal the average shown, as some beneficiaries moved from one county to another during this time frame.

SOCIOECONOMIC STATUS BY COUNTY

According to U.S. Census data from 2012, residents aged 65 and over in Atlantic County had a median household income of \$39,246. This was lower than the average income among all 10 counties (Figure 6).



Source: U.S. Census Bureau, American Community Survey (ACS), 2012 <http://www.census.gov/>.

PREVALENCE AND INCIDENCE

Using Medicare FFS claims data, eight behavioral health conditions were analyzed: depression, depression or proxy disorders, adjustment disorder, anxiety disorder, post-traumatic stress disorder (PTSD), substance abuse, alcohol or substance abuse, and suicide and intentional self-inflicted injury.

Claims data can underestimate the real incidence of depression in the population and individuals with depression could be diagnosed as having anxiety or adjustment disorders, as noted by the subject matter experts consulted for this project. Therefore, HQSI created a combination measure for depression (depression or proxy disorders) which includes beneficiaries who were reported for either depression, anxiety, or adjustment disorders.

The behavioral health data from January 1, 2011 to March 31, 2013 for these different measures were calculated to quantify disease occurrence:

1. Prevalence of the condition for the pre-Sandy time frame (Q1 2011 – Q3 2012, or 21 months)
2. Quarterly new incidence compared to prior year (Q1 2012 – Q1 2013)
3. The yearly prevalence of the condition with quarterly rolling trends to account for seasonal variation

Refer to Appendix A for measurement calculation and Appendix E for quarterly time frames and formulae.

Summary

Prevalence of the selected behavioral health conditions in the 21 months prior to Superstorm Sandy in the 10 counties is color coded with highest (red) and lowest (light blue) for each condition.

In the 21 months prior to Superstorm Sandy, Atlantic County had the highest prevalence rate of alcohol or substance abuse, substance abuse, and suicide and intentional self-inflicted injury among Medicare FFS beneficiaries in all 10 counties (Figure 7).


Figure 7. Prevalence of Selected Behavioral Health Conditions per 1,000 Medicare FFS Beneficiaries – Pre-Sandy (Q1 2011 – Q3 2012)

| County | Depression or Proxy Disorders | Depression | Anxiety Disorder | Adjustment Disorder | PTSD | Alcohol or Substance Abuse | Substance Abuse | Suicide and Intentional Self-Inflicted Injury |
|--------------------|-------------------------------|---------------|------------------|---------------------|-------------|----------------------------|-----------------|---|
| Atlantic | 269.50 | 171.36 | 163.83 | 45.32 | 7.13 | 60.37 | 34.67 | 9.24 |
| Bergen | 237.89 | 160.41 | 131.36 | 38.14 | 3.25 | 28.04 | 14.04 | 5.44 |
| Cape May | 273.48 | 169.64 | 168.71 | 36.62 | 6.74 | 55.75 | 28.95 | 6.22 |
| Essex | 242.98 | 158.79 | 124.32 | 54.16 | 4.92 | 56.59 | 33.80 | 7.02 |
| Hudson | 274.76 | 181.03 | 161.26 | 48.85 | 4.29 | 44.84 | 22.94 | 7.13 |
| Middlesex | 236.32 | 156.07 | 131.79 | 37.58 | 5.71 | 35.12 | 19.58 | 5.60 |
| Monmouth | 269.64 | 176.30 | 155.45 | 55.08 | 6.59 | 48.02 | 23.81 | 8.07 |
| Ocean | 272.03 | 173.97 | 170.82 | 42.95 | 7.48 | 50.61 | 30.52 | 8.47 |
| Somerset | 232.06 | 152.38 | 131.07 | 40.32 | 5.77 | 37.67 | 21.12 | 5.90 |
| Union | 225.96 | 148.14 | 126.46 | 31.27 | 3.32 | 34.96 | 17.45 | 5.12 |
| 10 counties | 251.86 | 164.75 | 144.53 | 43.45 | 5.36 | 43.26 | 23.75 | 6.80 |

■ Highest □ Lowest


Depression or Proxy Disorders

The prevalence rate of depression or proxy disorders in Atlantic County in the 21 months prior to Superstorm Sandy was 269.50 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 8).




The color-coded map of New Jersey depicts prevalence of depression or proxy disorders from high (red) to low (blue) in the 10 FEMA-declared disaster counties. This map shows that parts of Atlantic, along with Monmouth, Ocean, and Cape May Counties had a higher prevalence of depression or proxy disorders than the other counties (Figure 9).

FIGURE 9. PREVALENCE OF DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES (PRE-SANDY: Q1 2011- Q3 2012)




The color-coded map of Atlantic County depicts regional variation of prevalence of depression or proxy disorders from high (red) to low (blue) (Figure 10).

FIGURE 10. ATLANTIC COUNTY PREVALENCE OF DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES (PRE-SANDY: Q1 2011- Q3 2012)




For Q1 2013, there were 24.59 Medicare FFS beneficiaries newly diagnosed with depression or proxy disorders per 1,000 beneficiaries in Atlantic County compared to Q1 2012, which was 26.31 per 1,000 beneficiaries. This was higher than the quarterly rate among all 10 counties in Q1 2013 (Figure 11).




* Quarterly new incidences compared to prior year.


Comparing Q1 2013 to Q1 2012, there was a 6.5% relative decrease in new incidence of depression or proxy disorders in Atlantic County (Figure 12).



From Q2 2012 – Q1 2013, there was a 199.69 per 1,000 Medicare FFS beneficiaries yearly prevalence of depression or proxy disorders in Atlantic County compared to Q2 2011 – Q1 2012, which was 201.05 per 1,000 beneficiaries. Atlantic County’s yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 13).




Comparing Q2 2012 – Q1 2013 to Q2 2011 – Q1 2012, there was a 0.7% relative decrease in depression or proxy disorders in Atlantic County (Figure 14).




Depression

The prevalence rate of depression in Atlantic County in the 21 months prior to Superstorm Sandy was 171.36 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 15).




For Q1 2013, there were 16.12 Medicare FFS beneficiaries newly diagnosed with depression per 1,000 beneficiaries in Atlantic County compared to Q1 2012, which was 16.87 per 1,000 beneficiaries. This was higher than the quarterly rate among all 10 counties in Q1 2013 (Figure 16).




* Quarterly new incidences compared to prior year.


Comparing Q1 2013 to Q1 2012, there was a 4.4% relative decrease in new incidence of depression in Atlantic County (Figure 17).



From Q2 2012 – Q1 2013, there was a 121.06 per 1,000 Medicare FFS beneficiaries yearly prevalence of depression in Atlantic County compared to Q2 2011 – Q1 2012, which was 125.77 per 1,000 beneficiaries. Atlantic County’s yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 18).




Comparing Q2 2012 – Q1 2013 to Q2 2011 – Q1 2012, there was a 3.7% relative decrease in depression in Atlantic County (Figure 19).




Anxiety Disorders

The prevalence rate of anxiety disorders in Atlantic County in the 21 months prior to Superstorm Sandy was 163.83 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 20).




Adjustment Disorders

The prevalence rate of adjustment disorders in Atlantic County in the 21 months prior to Superstorm Sandy was 45.32 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 21).




Post-Traumatic Stress Disorder (PTSD)

The prevalence rate of PTSD in Atlantic County in the 21 months prior to Superstorm Sandy was 7.13 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 22).




For Q1 2013, there were 0.81 Medicare FFS beneficiaries newly diagnosed with PTSD per 1,000 beneficiaries in Atlantic County compared to Q1 2012, which was 0.87 per 1,000 beneficiaries. This was higher than the rate among all 10 counties in Q1 2013 (Figure 23).




* Quarterly new incidences compared to prior year.


Comparing Q1 2013 to Q1 2012, there was a 6.7% relative decrease in new incidence of PTSD in Atlantic County (Figure 24).



From Q2 2012 – Q1 2013, there was a 4.66 per 1,000 Medicare FFS beneficiaries yearly prevalence of PTSD in Atlantic County compared to Q2 2011 – Q1 2012, which was 5.07 per 1,000 beneficiaries. The yearly rate of PTSD with rolling quarters was higher than the rate among all 10 counties (Figure 25).




Comparing Q2 2012 – Q1 2013 to Q2 2011 – Q1 2012, there was an 8.1% relative decrease in PTSD in Atlantic County, the largest decrease among all 10 counties (Figure 26).




Alcohol or Substance Abuse

The alcohol or substance abuse measure includes Medicare FFS beneficiaries who were reported for either alcohol abuse or substance abuse.

The prevalence rate of alcohol or substance abuse in Atlantic County in the 21 months prior to Superstorm Sandy was 60.37 per 1,000 Medicare FFS beneficiaries. This was the highest prevalence rate among all 10 counties (Figure 27).




For Q1 2013, there were 6.72 Medicare FFS beneficiaries newly diagnosed with alcohol or substance abuse per 1,000 beneficiaries in Atlantic County compared to Q1 2012, which was 7.52 per 1,000 beneficiaries. This was higher than the quarterly rate among all 10 counties in Q1 2013 (Figure 28).




* Quarterly new incidences compared to prior year.


Comparing Q1 2013 to Q1 2012, there was a 10.7% relative decrease in new incidence of alcohol or substance abuse in Atlantic County (Figure 29).



From Q2 2012 – Q1 2013, there was a 39.05 per 1,000 Medicare FFS beneficiaries yearly prevalence of alcohol or substance abuse in Atlantic County compared to Q2 2011 – Q1 2012, which was 42.99 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 30).




Comparing Q2 2012 – Q1 2013 to Q2 2011 – Q1 2012, there was a 9.2% relative decrease in alcohol or substance abuse in Atlantic County, the largest decrease among all 10 counties (Figure 31).




Substance Abuse

The prevalence rate of substance abuse in Atlantic County in the 21 months prior to Superstorm Sandy was 34.67 per 1,000 Medicare FFS beneficiaries. This was the highest prevalence rate among all 10 counties (Figure 32).




For Q1 2013, there were 4.01 Medicare FFS beneficiaries newly diagnosed with substance abuse per 1,000 beneficiaries in Atlantic County compared to Q1 2012, which was 4.96 per 1,000 beneficiaries. This was higher than the quarterly rate among all 10 counties in Q1 2013 (Figure 33).




* Quarterly new incidences compared to prior year.


Comparing Q1 2013 to Q1 2012, there was a 19.2% relative decrease in new incidence of substance abuse in Atlantic County (Figure 34).



From Q2 2012 – Q1 2013, there was a 22.13 per 1,000 Medicare FFS beneficiaries yearly prevalence of substance abuse in Atlantic County compared to Q2 2011 – Q1 2012, which was 25.05 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 35).




Comparing Q2 2012 – Q1 2013 to Q2 2011 – Q1 2012, there was an 11.7% relative decrease in substance abuse in Atlantic County (Figure 36).




Suicide and Intentional Self-Inflicted Injury

The prevalence rate of suicide and intentional self-inflicted injury in Atlantic County in the 21 months prior to Superstorm Sandy was 9.24 per 1,000 Medicare FFS beneficiaries. This was the highest prevalence rate among all 10 counties (Figure 37).




For Q1 2013, there were 1.04 Medicare FFS beneficiaries newly diagnosed with suicide and intentional self-inflicted injury per 1,000 beneficiaries in Atlantic County, compared to Q1 2012, which was 1.03 per 1,000 beneficiaries. This is higher than the quarterly rate among all 10 counties in Q1 2013 (Figure 38).




* Quarterly new incidences compared to prior year.


Comparing Q1 2013 to Q1 2012, there was a 0.8% relative increase in new incidence of suicide and intentional self-inflicted injury in Atlantic County (Figure 39).



From Q2 2012 – Q1 2013, there was a 6.12 per 1,000 Medicare FFS beneficiaries yearly prevalence of suicide and intentional self-inflicted injury in Atlantic County compared to Q2 2011 – Q1 2012, which was 5.84 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 40).



Comparing Q2 2012 – Q1 2013 to Q2 2011 – Q1 2012, there was a 4.7% relative increase in suicide and intentional self-inflicted injury in Atlantic County (Figure 41).



RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS


To identify beneficiaries at risk of developing depression or proxy disorders, HQSI initially conducted a literature review on the potential risk factors for depression or proxy disorders. Previous studies suggested that psychosocial factors, biological factors, deteriorating physical functioning, and medication side effects could increase the risk of depression or proxy disorders.

Based on the literature review and running models using factors available through Medicare claims data, the top five risk factors for depression or proxy disorders were identified as: Alzheimer's disease and related disorders or senile dementia, sleep disturbance, substance or alcohol abuse or tobacco use, hip/pelvic fractures, and amputations (see Appendix B).

These risk factors were reported prior to the diagnosis of depression or proxy disorders thus indicating development of risk factors before diagnosis. The following figures show the prevalence rates for these five conditions in the 21 months prior to Superstorm Sandy.



TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS

The prevalence rate of beneficiaries with one or more of the top five risk factors for depression or proxy disorders in Atlantic County in the 21 months prior to Superstorm Sandy was 185.94 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 42).




The color-coded map of New Jersey depicts prevalence of any of the top five risk factors from high (red) to low (blue) in the 10 FEMA-declared disaster counties. This map shows that parts of Atlantic, along with Monmouth, Ocean, and Cape May Counties had higher prevalence of one or more of the top five risk factors for depression or proxy disorders (Figure 43).

FIGURE 43. PREVALENCE OF TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES (PRE-SANDY: Q1 2011 – Q3 2012)




The color-coded map of Atlantic County depicts regional variation of prevalence of any of the top five risk factors from high (red) to low (blue). Some areas, such as Atlantic City, Ventnor City, and Pleasantville had higher prevalence of one or more of the top five risk factors for depression or proxy disorders (Figure 44).

FIGURE 44. ATLANTIC COUNTY PREVALENCE OF TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES (PRE-SANDY: Q1 2011 – Q3 2012)




Alzheimer's Disease and Related Disorders or Senile Dementia

The prevalence rate of Alzheimer's disease and related disorders or senile dementia in Atlantic County in the 21 months prior to Superstorm Sandy was 42.31 per 1,000 Medicare FFS beneficiaries. This was the lowest prevalence rate among all 10 counties (Figure 45).




Sleep Disturbance

The prevalence rate of sleep disturbance in Atlantic County in the 21 months prior to Superstorm Sandy was 35.09 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 46).




Substance or Alcohol Abuse or Tobacco Use

The prevalence rate of substance or alcohol abuse or tobacco use in Atlantic County in the 21 months prior to Superstorm Sandy was 119.73 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 47).




Hip/Pelvic Fractures

The prevalence rate of hip/pelvic fractures in Atlantic County in the 21 months prior to Superstorm Sandy was 9.43 per 1,000 Medicare FFS beneficiaries. This was lower than the prevalence rate among all 10 counties (Figure 48).



Amputations

The prevalence rate of amputations in Atlantic County in the 21 months prior to Superstorm Sandy was 1.55 per 1,000 Medicare FFS beneficiaries. This was higher than the prevalence rate among all 10 counties (Figure 49).



—BEHAVIORAL HEALTH CONDITIONS—

This page left intentionally blank

OUTPATIENT BEHAVIORAL HEALTH SERVICES

Summary

HQSI analyzed five behavioral health assessment services and five behavioral health therapies. Utilization of outpatient health services is color coded with lowest (red) and highest (light blue).

The most frequently used behavioral health assessment service in Atlantic County was psychiatric diagnostic procedures (Figure 50).

Figure 50. Utilization of Outpatient Behavioral Health Services per 1,000 Medicare FFS Beneficiaries – Pre-Sandy Q1 2011 – Q3 2012 Assessments

| County | Depression Screening* | Diagnostic Psychological Tests | Health and Behavior Assessment/ Intervention | Neuropsychological Tests | Psychiatric Diagnostic Procedures |
|--------------------|-----------------------|--------------------------------|--|--------------------------|-----------------------------------|
| Atlantic | 1.12 | 4.59 | 1.46 | 12.04 | 90.99 |
| Bergen | 4.33 | 4.12 | 0.82 | 15.55 | 79.98 |
| Cape May | 0.65 | 2.71 | 0.94 | 8.98 | 72.68 |
| Essex | 0.83 | 7.81 | 0.97 | 12.77 | 88.05 |
| Hudson | 2.83 | 8.92 | 0.96 | 24.69 | 78.45 |
| Middlesex | 7.50 | 6.39 | 1.58 | 11.22 | 74.20 |
| Monmouth | 4.71 | 8.23 | 1.36 | 15.42 | 94.74 |
| Ocean | 9.49 | 4.39 | 1.03 | 14.83 | 82.92 |
| Somerset | 7.11 | 2.36 | 1.48 | 10.37 | 72.93 |
| Union | 3.02 | 2.63 | 1.33 | 10.60 | 71.56 |
| 10 counties | 4.81 | 5.53 | 1.16 | 14.18 | 81.61 |

* Depression screening rates are for Calendar Year 2012.



The most frequently used behavioral health therapy in Atlantic County was individual psychotherapy. Atlantic County had the highest utilization of group psychotherapy among all 10 counties (Figure 51).

Figure 51. Utilization of Outpatient Behavioral Health Services per 1,000 Medicare FFS Beneficiaries – Pre-Sandy Q1 2011 – Q3 2012
Therapies

| County | Biofeedback Therapy | Electroconvulsive Therapy | Family Psychotherapy | Group Psychotherapy | Individual Psychotherapy |
|-------------|---------------------|---------------------------|----------------------|---------------------|--------------------------|
| Atlantic | 1.20 | 0.40 | 4.02 | 7.83 | 69.00 |
| Bergen | 1.56 | 0.54 | 5.89 | 2.81 | 79.79 |
| Cape May | 0.89 | 0.98 | 1.64 | 2.01 | 50.28 |
| Essex | 0.88 | 0.63 | 6.04 | 3.23 | 78.83 |
| Hudson | 1.78 | 0.52 | 8.27 | 2.70 | 71.79 |
| Middlesex | 2.18 | 0.70 | 4.60 | 6.69 | 68.07 |
| Monmouth | 0.45 | 0.99 | 5.39 | 4.37 | 68.01 |
| Ocean | 0.80 | 0.84 | 3.76 | 2.11 | 61.53 |
| Somerset | 0.88 | 1.07 | 6.08 | 5.46 | 73.16 |
| Union | 0.84 | 0.54 | 3.25 | 3.29 | 55.35 |
| 10 counties | 1.19 | 0.70 | 5.09 | 3.88 | 69.31 |




Assessments

Depression Screening

One of the long-term goals of this project is to increase the awareness and use of Medicare-covered depression screening among at-risk Medicare FFS beneficiaries residing in the 10 counties during Superstorm Sandy. The depression screening utilization rates have been low in all 10 counties.


Beginning January 2012, depression screening became a Medicare-covered service. According to the CMS Screening for Depression Booklet, Medicare Part B covers an annual screening for depression of 15 minutes in length for Medicare FFS beneficiaries in primary care settings when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up.

The rate of depression screening in Atlantic County for calendar year 2012 was 1.12 per 1,000 Medicare FFS beneficiaries. This was lower than the rate among all 10 counties (Figure 52).



The color-coded map of New Jersey depicts the use of depression screening from low (red) to high (blue) in the 10 FEMA-declared disaster counties (Figure 53).

FIGURE 53. DEPRESSION SCREENING PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES (CALENDAR YEAR 2012)



The color-coded map of Atlantic County depicts regional variation in the rates of the use of the depression screening benefit from low (red) to high (blue) (Figure 54).

FIGURE 54. ATLANTIC COUNTY DEPRESSION SCREENING PER 1,000 MEDICARE FFS BENEFICIARIES (CALENDAR YEAR 2012)






Figure 55 shows five quarters of data to reflect trending in the use of the depression screening benefit per 1,000 Medicare FFS beneficiaries in Atlantic County and in the 10 FEMA-declared disaster counties.



Diagnostic Psychological Tests

According to the CMS Mental Health Services Billing Guide, psychological testing includes psychodiagnostic assessment of emotionality, intellectual abilities, personality, and psychopathology (e.g., Minnesota Multiphasic Personality Inventory, Rorschach, or Wechsler Adult Intelligence Scale).


The rate of psychological testing in Atlantic County in the 21 months prior to Superstorm Sandy was 4.59 per 1,000 Medicare FFS beneficiaries. This was lower than the rate among all 10 counties (Figure 56).



Health and Behavior Assessment/Intervention

According to the CMS Mental Health Services Billing Guide, health and behavior assessments are used to identify the psychological, behavioral, emotional, cognitive, and social factors important to the prevention, treatment, or management of physical health problems.

The rate of health and behavior assessment in Atlantic County in the 21 months prior to Superstorm Sandy was 1.46 per 1,000 Medicare FFS beneficiaries. This was higher than the rate among all 10 counties (Figure 57).



Neuropsychological Tests

According to the CMS Mental Health Services Billing Guide, neuropsychological tests are evaluations designed to determine the functional consequences of known or suspected brain injury through testing of the neurocognitive domains responsible for language, perception, memory, learning, problem solving, and adaptation.

The rate of neuropsychological testing in Atlantic County in the 21 months prior to Superstorm Sandy was 12.04 per 1,000 Medicare FFS beneficiaries. This was lower than the rate among all 10 counties (Figure 58).





Figure 59 shows nine quarters of data to reflect trending in the use of neuropsychological testing per 1,000 Medicare FFS beneficiaries in Atlantic County and in the 10 FEMA-declared disaster counties.



Psychiatric Diagnostic Procedures

According to the CMS Mental Health Services Billing Guide, psychiatric diagnostic evaluation is an integrated biopsychosocial assessment, including history, mental status, and recommendations. The evaluation may include communication with family or other sources and review of diagnostic studies.

The utilization rate of psychiatric diagnostic procedures in Atlantic County in the 21 months prior to Superstorm Sandy was 90.99 per 1,000 Medicare FFS beneficiaries. This was higher than the rate among all 10 counties (Figure 60).





Figure 61 shows nine quarters of data to reflect trending in the use of psychiatric diagnostic procedures per 1,000 Medicare FFS beneficiaries in Atlantic County and in the 10 FEMA-declared disaster counties.



Therapies

Individual Psychotherapy

According to the CMS Mental Health Services Billing Guide, individual psychotherapy is the treatment of mental illness and behavioral disturbances where the physician or other qualified health professional attempts to alleviate the emotional disturbances, reverse or change maladaptive patterns of behavior, and encourage personality growth and development. This is done through the use of definitive therapeutic communication.

The utilization rate of individual psychotherapy in Atlantic County in the 21 months prior to Superstorm Sandy was 69.00 per 1,000 Medicare FFS beneficiaries. This was lower than the rate among all 10 counties (Figure 62).





Figure 63 shows nine quarters of data to reflect trending in the use of individual psychotherapy per 1,000 Medicare FFS beneficiaries in Atlantic County and in the 10 FEMA-declared disaster counties.




Family Psychotherapy

According to the CMS Mental Health Services Billing Guide, family psychotherapy describes the treatment of the family unit when maladaptive behaviors of family members are exacerbating the beneficiary’s mental illness or interfering with treatment. It can also be used to assist the family in addressing the maladaptive behaviors of the patient and improve treatment compliance.

The utilization rate of family psychotherapy was 4.02 per 1,000 Medicare FFS beneficiaries in Atlantic County in the 21 months prior to Superstorm Sandy. This was lower than the rate among all 10 counties (Figure 64).

Due to these low numbers, no quarterly trending data has been provided for this therapy.




Group Psychotherapy

According to the CMS Mental Health Services Billing Guide, group psychotherapy is a form of treatment where a selected group of patients are guided by a licensed psychotherapist for the purpose of helping to change maladaptive patterns which interfere with social functioning and are associated with a diagnosable psychiatric illness.

The utilization rate of group psychotherapy was 7.83 per 1,000 Medicare FFS beneficiaries in Atlantic County in the 21 months prior to Superstorm Sandy. This was the highest rate among all 10 counties (Figure 65).

Due to these low numbers, no quarterly trending data has been provided for this therapy.




Electroconvulsive Therapy

According to the CMS Mental Health Services Billing Guide, electroconvulsive therapy (ECT) is the application of electric current to the brain through scalp electrodes to induce a single seizure to produce a therapeutic effect. It is used primarily to treat major depressive disorder when antidepressant medication should not be used because it may be harmful to the patient. This type of therapy can be used for certain other clinical conditions as well.

The utilization rate of ECT was 0.40 per 1,000 Medicare FFS beneficiaries in Atlantic County in the 21 months prior to Superstorm Sandy. This was the lowest rate among all 10 counties (Figure 66).

Due to these low numbers, no quarterly trending data has been provided for this therapy.




Biofeedback Therapy

According to the CMS Mental Health Services Billing Guide, biofeedback therapy provides visual, auditory, or other evidence of the status of certain body functions so that a person can exert voluntary control over those functions, and thereby alleviate an abnormal bodily condition.

The utilization rate of biofeedback therapy rate was 1.20 per 1,000 Medicare FFS beneficiaries in Atlantic County in the 21 months prior to Superstorm Sandy. This was higher than the rate among all 10 counties (Figure 67).

Due to these low numbers, no quarterly trending data has been provided for this therapy.



INPATIENT SERVICES

Summary

Utilization of inpatient health services per 1,000 Medicare FFS beneficiaries in the 21 months prior to Superstorm Sandy in the 10 counties is color coded with highest (red) and lowest (light blue) for each measure. These additional services were analyzed because beneficiaries with underlying behavioral health issues may seek non-behavioral health services.

Inpatient services included four measures of service utilization: inpatient psychiatric facilities, acute care hospital admissions, observation stays with or without a subsequent hospital admission, and emergency department visits with or without a subsequent hospital admission. The second set of measures is tied to utilization of services within 30 days of an acute care episode, often used as proxy indicators of care coordination, and include all-cause 30-day hospital readmissions, observation stays within 30 days of discharge, and emergency department visits within 30 days of discharge.

Atlantic County had the highest rate of acute care hospital admissions and emergency department visits among all 10 counties (Figure 68).

Figure 68. Utilization of Inpatient Health Services per 1,000 Medicare FFS Beneficiaries – Pre-Sandy (Q1 2011-Q3 2012)

| County | Psychiatric Hospital Admissions | Acute Care Hospital Admissions | Observation Stays | Emergency Department Visits |
|--------------------|---------------------------------|--------------------------------|-------------------|-----------------------------|
| Atlantic | 10.58 | 642.49 | 25.54 | 510.52 |
| Bergen | 13.31 | 514.96 | 11.18 | 390.06 |
| Cape May | 7.81 | 639.80 | 8.00 | 474.60 |
| Essex | 14.35 | 625.28 | 53.49 | 502.88 |
| Hudson | 14.25 | 611.20 | 34.01 | 459.27 |
| Middlesex | 10.33 | 535.00 | 25.09 | 427.85 |
| Monmouth | 13.96 | 575.84 | 8.23 | 448.58 |
| Ocean | 12.34 | 602.84 | 7.82 | 455.17 |
| Somerset | 12.58 | 502.79 | 8.34 | 392.45 |
| Union | 11.77 | 500.74 | 12.13 | 395.58 |
| 10 counties | 12.56 | 567.94 | 19.45 | 440.57 |



Atlantic County had the highest rate of emergency department visits within 30 days of discharge among all 10 counties (Figure 69).


Figure 69. Utilization of Inpatient Health Services per 1,000 Medicare FFS Beneficiaries Within 30 Days of Discharge Pre-Sandy (Q1 2011-Q3 2012)

| County | 30-Day Hospital Readmissions | Observation Stays | Emergency Department Visits |
|-------------|------------------------------|-------------------|-----------------------------|
| Atlantic | 129.80 | 14.35 | 167.45 |
| Bergen | 96.03 | 7.45 | 112.75 |
| Cape May | 117.17 | 10.66 | 150.19 |
| Essex | 142.86 | 21.39 | 164.37 |
| Hudson | 139.88 | 19.76 | 155.26 |
| Middlesex | 104.29 | 12.81 | 124.80 |
| Monmouth | 109.54 | 9.46 | 132.57 |
| Ocean | 115.31 | 11.52 | 143.97 |
| Somerset | 93.09 | 7.72 | 112.03 |
| Union | 92.61 | 9.08 | 110.27 |
| 10 counties | 112.70 | 12.25 | 134.83 |


■ Highest □ Lowest

Psychiatric Hospital Admissions

Standalone psychiatric hospitals or distinct part psychiatric units in acute care hospitals in Atlantic County in the 21 months prior to Superstorm Sandy had an admissions rate of 10.58 per 1,000 Medicare FFS beneficiaries (Figure 70).



From Q2 2012 – Q1 2013, there were 5.91 per 1,000 Medicare FFS beneficiaries admitted to psychiatric hospitals in Atlantic County compared to Q2 2011 – Q1 2012, which was 7.97 per 1,000 beneficiaries. The yearly rate of psychiatric hospital admissions with rolling quarters was lower than the rate among all 10 counties (Figure 71).




Acute Care Hospitals


Admissions

The following data shows all-cause utilization measures and includes all Medicare FFS beneficiaries, not just beneficiaries with behavioral health conditions. The data are ranked from lowest utilization (top) to the highest utilization (bottom).

The rate of hospital admissions in Atlantic County in the 21 months prior to Superstorm Sandy was 642.49 per 1,000 Medicare FFS beneficiaries. This was the highest admissions rate among all 10 counties (Figure 72).




From Q2 2012 – Q1 2013, the hospital admissions rate was 351.85 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 365.38 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 73).




Observation Stays

According to the U.S. Department of Health and Human Services, observation stays are short-term treatments and assessments provided to outpatients to determine whether Medicare FFS beneficiaries require further treatment as inpatients or can be discharged.

The rate of observation stays in Atlantic County in the 21 months prior to Superstorm Sandy was 25.54 per 1,000 Medicare FFS beneficiaries. This was higher than the rate among all 10 counties (Figure 74).




From Q2 2012 – Q1 2013, the rate of observation stays was 16.92 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 15.60 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 75).




Emergency Department Visits

The rate of emergency department visits in Atlantic County in the 21 months prior to Superstorm Sandy was 510.52 per 1,000 Medicare FFS beneficiaries. This was the highest rate among all 10 counties (Figure 76).




From Q2 2012 – Q1 2013, the rate of emergency department visits was 282.88 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 288.74 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 77).




30-Day Hospital Readmissions

The rate of 30-day hospital readmissions in Atlantic County in the 21 months prior to Superstorm Sandy was 129.80 per 1,000 Medicare FFS beneficiaries. This was higher than the rate among all 10 counties (Figure 78).




From Q2 2012 – Q1 2013, the rate of 30-day hospital readmissions was 68.73 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 74.12 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 79).




Observation Stays Within 30 Days of Discharge

The rate of observation stays within 30 days of discharge in Atlantic County in the 21 months prior to Superstorm Sandy was 14.35 per 1,000 Medicare FFS beneficiaries. This was higher than the rate among all 10 counties (Figure 80).




From Q2 2012 – Q1 2013, the rate of observation stays within 30 days of discharge was 7.60 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 9.33 per 1,000 beneficiaries. The yearly rate of observation stays within 30 days of discharge with rolling quarters was higher than the rate among all 10 counties (Figure 81).




Emergency Department Visits Within 30 Days of Discharge

The rate of emergency department visits within 30 days of discharge in Atlantic County in the 21 months prior to Superstorm Sandy was 167.45 per 1,000 Medicare FFS beneficiaries. This was the highest rate among all 10 counties (Figure 82).



From Q2 2012 – Q1 2013, the rate of emergency department visits within 30 days of discharge was 90.49 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 94.95 per 1,000 beneficiaries. The yearly rate of emergency department visits within 30 days of discharge with rolling quarters was higher than the rate among all 10 counties (Figure 83).



Other Settings

Summary

This profile also looks at the utilization of home health agency, skilled nursing facility, hospice, and medical rehabilitation services. These additional services were analyzed because Medicare FFS beneficiaries with underlying behavioral health issues may seek these non-behavioral health services. Utilization of health services per 1,000 Medicare FFS beneficiaries for these settings in the 21 months prior to Superstorm Sandy in the 10 counties is color coded with highest (red) and lowest (light blue) for each measure.

Atlantic County had the lowest utilization rate of skilled nursing facility services among all 10 counties (Figure 84).


**Figure 84. Utilization of Health Services
per 1,000 Medicare FFS Beneficiaries – Pre-Sandy (Q1 2011-Q3 2012)**

| County | Home Health Agency Services | Skilled Nursing Facility Services | Hospice Services | Medical Rehabilitation Services |
|--------------------|-----------------------------|-----------------------------------|------------------|---------------------------------|
| Atlantic | 144.57 | 99.09 | 44.90 | 22.65 |
| Bergen | 134.21 | 103.49 | 34.63 | 18.52 |
| Cape May | 141.72 | 106.74 | 46.96 | 13.94 |
| Essex | 121.71 | 118.04 | 31.68 | 15.32 |
| Hudson | 138.52 | 110.10 | 27.96 | 13.43 |
| Middlesex | 124.56 | 107.09 | 34.23 | 15.01 |
| Monmouth | 146.06 | 112.50 | 48.77 | 26.61 |
| Ocean | 153.02 | 118.04 | 49.43 | 36.49 |
| Somerset | 125.70 | 103.72 | 39.73 | 16.60 |
| Union | 126.23 | 109.34 | 33.87 | 15.52 |
| 10 counties | 135.89 | 109.87 | 38.86 | 20.79 |




Home Health Agency Services

The rate of home health agency use in Atlantic County in the 21 months prior to Superstorm Sandy was 144.57 per 1,000 Medicare FFS beneficiaries. This was higher than the rate among all 10 counties (Figure 85).




From Q2 2012 – Q1 2013, the utilization rate of home health agency services was 93.76 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 99.78 per 1,000 beneficiaries. The yearly rate of home health agency use with rolling quarters was higher than the rate among all 10 counties (Figure 86).




Skilled Nursing Facility Services

The rate of skilled nursing facility services in Atlantic County in the 21 months prior to Superstorm Sandy was 99.09 per 1,000 Medicare FFS beneficiaries. This was the lowest utilization rate among all 10 counties (Figure 87).




From Q2 2012 – Q1 2013, the utilization rate of skilled nursing facility services was 62.55 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 64.27 per 1,000 beneficiaries. The yearly rate of skilled nursing facility use with rolling quarters was lower than the rate among all 10 counties (Figure 88).




Hospice Services

The rate of hospice use in Atlantic County in the 21 months prior to Superstorm Sandy was 44.90 per 1,000 Medicare FFS beneficiaries. This was higher than the rate among all 10 counties (Figure 89).




From Q2 2012 – Q1 2013, the rate of hospice use was 29.52 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 28.04 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 90).




Medical Rehabilitation Services

The rate of medical rehabilitation use in Atlantic County in the 21 months prior to Superstorm Sandy was 22.65 per 1,000 Medicare FFS beneficiaries. This was higher than the rate among all 10 counties (Figure 91).



From Q2 2012 – Q1 2013, the rate of medical rehabilitation use was 13.09 per 1,000 Medicare FFS beneficiaries in Atlantic County compared to Q2 2011 – Q1 2012, which was 13.93 per 1,000 beneficiaries. The yearly rate with rolling quarters was higher than the rate among all 10 counties (Figure 92).



APPENDIX A: BEHAVIORAL HEALTH CONDITIONS

Documentation and Technical Notes

The following defines the study population, the time frames, and the exclusion and inclusion criteria:

Data Source

- New Jersey Medicare Part A and Part B FFS claims data and denominator file

Reference Time Period

- Prevalence of the condition for the pre-Sandy time frame (Q1 2011- Q3 2012 or 21 months)
- Yearly prevalence of the condition with quarterly rolling (Q1 2011-Q1 2013)
- Quarterly new incidence of conditions that were not existent (not reported) in the prior year

Denominator

- All Medicare beneficiaries who were in CMS denominator file during measurement time frame
- With FFS coverage AND eligible enrollment in FFS days/total measurement days > 0

Numerator

- Unique beneficiaries with disease-specific inpatient OR outpatient claim during the time frame
- CCW and AHRQ disease diagnosis code match (ICD-9-CM codes) Part A dgns_cd_1-25 and dgns_e_cd_1-3; Match Part B dgns_cd_1_12

Exclusions

- HMO coverage period
- Age <18 or >= 110
- Eligible FFS days/total measurement days = 0

Resources

More information on the classification codes, requirements, and processing of the behavioral health conditions highlighted in this profile can be located at the following links:

- https://www.ccwdata.org/cs/groups/public/documents/document/clin_cond_algo_req_proc.pdf
- <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>

APPENDIX A

The following table shows the ICD-9-CM codes for the eight behavioral health conditions:

| Behavioral Health Conditions | Numerator: Valid ICD-9-CM Codes |
|---|---|
| Depression or Proxy Disorders (Depression, Anxiety Disorders or Adjustment Disorders) | 29384, 29620, 29621, 29622, 29623, 29624, 29625, 29626, 29630, 29631, 29632, 29633, 29634, 29635, 29636, 30000, 30001, 30002, 30009, 30010, 30020, 30021, 30022, 30023, 30029, 3003, 3004, 3005, 30089, 3009, 3080, 3081, 3082, 3083, 3084, 3089, 3090, 3091, 30922, 30923, 30924, 30928, 30929, 3093, 3094, 30981, 30982, 30983, 30989, 3099, 311, 3130, 3131, 31321, 31322, 3133, 31382, 31383, V790 |
| Depression | 29620, 29621, 29622, 29623, 29624, 29625, 29626, 29630, 29631, 29632, 29633, 29634, 29635, 29636, 3004, 311, V790 |
| Anxiety Disorders | 29384, 30000, 30001, 30002, 30009, 30010, 30020, 30021, 30022, 30023, 30029, 3003, , 3005, 30089, 3009, 3080, 3081, 3082, 3083, 3084, 3089, 3130, 3131, 31321, 31322, 3133, 31382, 31383 |
| Adjustment Disorders | 3090, 3091, 30922, 30923, 30924, 30928, 30929, 3093, 3094, 30981, 30982, 30983, 30989, 3099 |
| Post-Traumatic Stress Disorder (PTSD) | 30981 |
| Alcohol or Substance Abuse | 2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76072, 76073, 76075, 7795, 96500, 96501, 96502, 96509, V6542 Alcohol Abuse: 2910, 2911, 2912, 2913, 2914, 2915, 2918, 29181, 29182, 29189, 2919, 30300, 30301, 30302, 30303, 30390, 30391, 30392, 30393, 30500, 30501, 30502, 30503, 76071, 9800 |
| Substance Abuse | 2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76072, 76073, 76075, 7795, 96500, 96501, 96502, 96509, V6542 |
| Suicide and Intentional Self-Inflicted Injury | E9500, E9501, E9502, E9503, E9504, E9505, E9506, E9507, E9508, E9509, E9510, E9511, E9518, E9520, E9521, E9528, E9529, E9530, E9531, E9538, E9539, E954, E9550, E9551, E9552, E9553, E9554, E9555, E9556, E9557, E9559, E956, E9570, E9571, E9572, E9579, E9580, E9581, E9582, E9583, E9584, E9585, E9586, E9587, E9588, E9589, E959, V6284 |

APPENDIX B: RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS

Documentation and Technical Notes

The following defines the study population, the time frame, the exclusion and inclusion criteria, and the literature review references:

Data Source

- New Jersey Medicare Part A and Part B FFS claims data and denominator file

Reference Time Period

- Prevalence of the condition for the pre-Sandy time frame (January 2011 – September 2012 or 21 months)

Denominator

- All Medicare beneficiaries who were in CMS denominator file during measurement time frame
- With FFS coverage AND eligible enrollment in FFS days/total measurement days > 0

Numerator

- Unique beneficiaries with disease-specific inpatient OR outpatient claim during the time frame
- CCW and AHRQ disease diagnosis code match (ICD-9-CM codes) Part A dgns_cd_1-25 and dgns_e_cd_1-3; Match Part B dgns_cd_1_12

Exclusions

- HMO coverage period
- Age <18 or >= 110
- Eligible FFS days/total measurement days = 0

Model

- Logistic Regression Models were used to determine the top five risk factors with the highest Odds Ratios (OR) ($p < 0.001$)

Resources

More information on the classification codes, requirements, and processing of the combination measure of depression or proxy disorders which includes beneficiaries reported for either depression, anxiety, or adjustment disorders can be located at the following links:

- https://www.ccwdata.org/cs/groups/public/documents/document/clin_cond_algo_req_proc.pdf
- <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>

Literature Review References for Risk Factors for Depression or Proxy Disorders

National Alliance on Mental Illness. Information Helpline: Depression in Older Persons Fact Sheet [Internet]. Arlington (VA): National Alliance on Mental Illness; 2009 Oct [cited 2013 Sep 17]. Available from: <http://www.nami.org/Template.cfm?Section=Helpline1&Template=/ContentManagement/ContentDisplay.cfm&ContentID=144039>

National Institute of Mental Health. Depression: Causes and Risk Factors [Internet]. Bethesda (MD): National Institute of Mental Health; 2013 Jul [cited 2013 Sep 17]. Available from: <http://nihseniorhealth.gov/depression/causesandriskfactors/01.html>

Centers for Disease Control and Prevention and National Association of Chronic Disease Directors. The State of Mental Health and Aging in America [Internet]. Atlanta (GA): National Association of Chronic Disease Directors, 2008 [cited 2013 Sep 19]. 11 p. Available from: http://www.cdc.gov/aging/pdf/mental_health.pdf

Jacques L, Jensen T, Schafer J, Caplan S, Schott L. Final Coverage Decision Memorandum for Screening for Depression in Adults [Internet]. Baltimore (MD): Centers for Medicare & Medicaid Services; 2011 Oct 14 [cited 2013 Sep 18]. 42 p. Available from: <http://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=251>

Thakur M, Blazer DG. Depression in long-term care. Journal of the American Medical Directors Association [Internet]. 2008 Feb [cited 2013 Sep 19];9(2):82-87. Available from: <http://www.amda.com/tools/clinical/depression/DepressioninLongTermCare.pdf>

Sozeri-Varma G. Depression in the elderly: clinical features and risk factors. Aging and Disease [Internet]. 2012 Dec [cited 2013 Sep 18];3(6):465-471. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3522513/>

Qian J, Simoni-Wastila L, Rattinger GB, Lehmann S, Langenberg P, et al. Associations of depression diagnosis and antidepressant treatment with mortality among young and disabled Medicare beneficiaries with COPD. General Hospital Psychiatry. 2013 Jul 18 [cited 2013 Sep 22]; 35(6):612-618.

Shao W, Ahmad R, Khutoryansky N, Aagren M, Bouchard J. Evidence supporting an association between hypoglycemic events and depression. Current Medical Research and Opinion. 2013 Sep 23 [cited 2013 Sep 22]: 1-7.

Substance Abuse and Mental Health Services Administration. The Treatment of Depression in Older Adults: Depression and Older Adults: Key Issues [Internet]. Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Services Administration, U.S. Department of Health and Human Services, 2011 [cited 2013 Sep 24]. HHS Pub. No. SMA-11-4631. 24 p. Available from: <http://store.samhsa.gov/shin/content/SMA11-4631CD-DVD/SMA11-4631CD-DVD-KeyIssues.pdf>

Himelhoch S, Weller WE, Wu AW, Anderson GF, Cooper LA. Chronic medical illness, depression, and use of acute medical services among Medicare beneficiaries. Medical Care. 2004 Jun [cited 2013 Sep 25];42(6):512-521.

Mohile SG, Fan L, Reeve E, Jean-Pierre P, Mustian K, et al. Association of cancer with geriatric syndromes in older Medicare beneficiaries. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* [Internet]. 2011 Apr 10 [cited 2013 Sep 25];29(11):1458-1464. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3082984/>

Jayadevappa R, Malkowicz SB, Chhatre S, Johnson JC, Gallo JJ. The burden of depression in prostate cancer. *Psycho-oncology*. 2012 Dec [cited 2013 Sep 26];21(12):1338-1345.

Missouri Department of Mental Health. CPS Facts: Depression and Older Adults [Internet]. Jefferson City(MO): Missouri Department of Mental Health, [date unknown, cited 2013 Sep 26], 2 p. Available from: <http://dmh.mo.gov/docs/mentalillness/elderlydepress.pdf>

Oregon State University, Washington State University, University of Idaho. Depression in Later Life: Recognition and Treatment [Internet]. Corvallis(OR): Pacific Northwest Extension Publication; 2004 Jul [Published April 1990; revised July 2000; cited 2013 Sep 29]; 32 p. Available from: <http://ir.library.oregonstate.edu/xmlui/bitstream/handle/1957/20713/pnw347.pdf>

Cole MG, Dendukuri N. Risk factors for depression among elderly community subjects: a systematic review and meta-analysis. *American Journal of Psychiatry* [Internet]. 2003 Jun [cited 2013 Sep 29]; 160(6):1147-1156. Available from: <http://ajp.psychiatryonline.org/article.aspx?articleid=176272>

Kohn R, Levav I, Garcia ID, Machuca ME, Tamashiro R. Prevalence, risk factors and aging vulnerability for psychopathology following a natural disaster in a developing country. *International Journal of Geriatric Psychiatry*. 2005 Sep [cited 2013 Sep 29];20(9):835-841.

Pietrzak RH, Southwick SM, Tracy M, Galea S, Norris FH. Posttraumatic stress disorder, depression, and perceived needs for psychological care in older persons affected by Hurricane Ike. *Journal of Affective Disorders* [Internet]. 2012 Apr [cited 2013 Sep 30];138(1-2):96-103. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3306486/>

Oriol W. Psychosocial Issues for Older Adults in Disasters [Internet]. Washington (DC): Emergency Services and Disaster Relief Branch, Center for Mental Health Services (CMHS), Substance Abuse and Mental Health Services Administration; 1999 [cited 2013 Sep 30]; DHHS Publication No. ESDRB SMA 99-3323. 79 p. Available from: <http://store.samhsa.gov/shin/content/SMA99-3323/SMA99-3323.pdf>

O'Connor EA, Whitlock EP, Gaynes B, Beil TL. Screening for Depression in Adults and Older Adults in Primary Care: An Updated Systematic Review. [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2009 Dec [cited 2013 Sept 30]. 167 p. (Evidence Synthesis No. 75. AHRQ Publication No. 10-05143-EF-1). Available from: <http://www.ncbi.nlm.nih.gov/books/NBK36403/pdf/TOC.pdf>

Noyes K, Liu H, Lyness JM, Friedman B. Medicare beneficiaries with depression: comparing diagnoses in claims data with the results of screening. *Psychiatric Services* [Internet]. 2011 Oct [cited 2013 Sep 30];62(10):1159-1166. Available from: http://ps.psychiatryonline.org/data/Journals/PSS/4336/pss6210_1159.pdf

APPENDIX B

The following table shows the ICD-9-CM codes for the top five risk factors for depression or proxy disorders:

| Top Five Risk Factors for Depression or Proxy Disorders* | Numerator: Valid ICD-9-CM Codes |
|--|---|
| Alzheimer's Disease and Related Disorders or Senile Dementia | 3311, 33111, 33119, 3312, 3317, 2900, 29010, 29011, 29012, 29013, 29020, 29021, 2903, 29040, 29041, 29042, 29043, 2940, 2941, 29410, 29411, 2948, 797 |
| Sleep Disturbance | 04672, 29182, 29285, 30740, 30741, 30742, 30748, 30749, 32700, 32701, 32702, 32709, 78050, 78051, 78052, 78059 |
| Substance or Alcohol Abuse or Tobacco Use | 2910, 2911, 2912, 2913, 2914, 2915, 2918, 29181, 29182, 29189, 2919, 2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30300, 30301, 30302, 30303, 30390, 30391, 30392, 30393, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30500, 30501, 30502, 30503, 3051, 30510, 30511, 30512, 30513, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 33392, 3575, 4255, 5353, 53530, 53531, 5710, 5711, 5712, 5713, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76071, 76072, 76073, 76075, 7795, 7903, 96500, 96501, 96502, 96509, 9800, V110, V111, V112, V113, V114, V118, V119, V154, V1541, V1542, V1549, V1582, V6285, V6542, V663, V701, V702, V7101, V7102, V7109, V790, V791, V792, V793, V798, V799 |
| Hip/Pelvic Fractures | 73314, 73315, 73396, 73397, 73398, 8080, 8081, 8082, 8083, 80841, 80842, 80843, 80849, 80851, 80852, 80853, 80859, 8088, 8089, 82000, 82001, 82002, 82003, 82009, 82010, 82011, 82012, 82013, 82019, 82020, 82021, 82022, 82030, 82031, 82032, 8208, 8209 |
| Amputations | 8870, 8871, 8872, 8873, 8874, 8875, 8876, 8877, 8960, 8961, 8962, 8963, 8970, 8971, 8972, 8973, 8974, 8975, 8976, 8977, 9059, 99760, 99761, 99762, 99769 |

* Other risk factors for depression or proxy disorders analyzed include Acute Myocardial Infarction (AMI), Stroke/Transient Ischemic Attack, Coronary Artery Bypass Graft Surgery (CABG), Parkinson's Disease, Chronic Obstructive Pulmonary Disease and Bronchiectasis (COPD), Diabetes, Chronic Kidney Disease, Rheumatoid Arthritis/Osteoarthritis (RA/OA), Macular Degeneration, Disability, History of Cancer, Heart Failure, and Acquired Hypothyroidism.

APPENDIX C: UTILIZATION OF OUTPATIENT MENTAL HEALTH SERVICES

Documentation and Technical Notes

The following defines the study population, the time frame, and the exclusion and inclusion criteria:

Data Source

- New Jersey Medicare Part A and Part B FFS claims data and denominator file

Reference Time Period

- Utilization during pre-Sandy time frame (January 2011 – September 2012 or 21 months)
- Depression Screening: Calendar Year (CY) 2012
- Quarterly utilization (January 2011 – March 2013 or nine quarters)

Denominator

- All Medicare beneficiaries who were in CMS denominator file during measurement time frame
- With FFS coverage AND eligible enrollment in FFS days/total measurement days > 0

Numerator

Unique beneficiaries with specific outpatient mental health service claims

Exclusions

- HMO coverage period
- Age <18 or >= 110
- Eligible FFS days/total measurement days =0

Resources

More information on the definitions and uses of the outpatient mental health services highlighted in this profile can be located at <http://www.medicarenhic.com/providers/pubs/REF-EDO-0012MentalHealthBillingGuide2013.pdf>.

APPENDIX C

The following table shows the CPT/HCPCS codes for the outpatient mental health services:

| Mental Health Services | Numerator: CPT/HCPCS Codes |
|---|--|
| Assessments | |
| Depression Screening | G0444 |
| Diagnostic Psychological Tests | 96101, 96102, 96103, 96105, 96110, 96111 |
| Health and Behavior Assessment/Intervention | 96150, 96151, 96152, 96153, 96154, 96155 |
| Neuropsychological Tests | 96116, 96118, 96119, 96120 |
| Psychiatric Diagnostic Procedures | 90801, 90802, 90791, 90792 |
| Therapies | |
| Individual Psychotherapy | 90804, 90805, 90832, 90833, 90806, 90807, 90834, 90836, 90808, 90809, 90810, 90811, 90812, 90813, 90814, 90815, 90816, 90817, 90818, 90819, 90821, 90822, 90823, 90824, 90826, 90827, 90828, 90829, 90837, 90838, 90839, 90840 |
| Family Psychotherapy | 90846, 90847 |
| Group Psychotherapy | 90849, 90853, 90857 |
| Electroconvulsive Therapy | 90870 |
| Biofeedback Therapy | 90901, 90911 |

APPENDIX D: UTILIZATION OF SERVICES – INPATIENT AND OTHER SETTINGS

Documentation and Technical Notes

The following defines the study population, the time frame, and the exclusion and inclusion criteria:

Data Source

New Jersey Medicare Part A and Part B FFS claims data and denominator file

Reference Time Period

- Utilization during pre-Sandy time frame (January 2011 – September 2012 or 21 months)
- Quarterly utilization of services (January 2011 – March 2013 or nine quarters)
- Yearly utilization of services with quarterly rolling (January 2011 – March 2013)

Denominator

- All Medicare beneficiaries who were in CMS denominator file during measurement time frame
- With FFS coverage AND eligible enrollment in FFS days/total measurement days > 0

Exclusions

- HMO coverage period
- Age < 18 or >= 110
- Eligible FFS days/total measurement days = 0

Utilization Measure

Refer to Appendix E.

Numerator

| Utilization Measure Description | Numerator |
|--|---|
| Psychiatric Hospital Admissions | Number of eligible beneficiaries with at least one psychiatric hospital admission claim |
| Acute Care Hospital Admissions | Number of acute care hospital admissions |
| Observation Stays | Number of observation stays |
| Emergency Department Visits | Number of emergency department visits |
| 30-Day Hospital Readmissions | Number of 30-day hospital readmissions |
| Observation Stays Within 30 Days of Hospital Discharge | Number of observation stays within 30 days of hospital discharge |
| Emergency Department Visits Within 30 Days of Hospital Discharge | Number of emergency department visits within 30 days of hospital discharge |
| Home Health Agency Services | Number of eligible beneficiaries with at least one home health agency claim |
| Skilled Nursing Facility Services | Number of eligible beneficiaries with at least one skilled nursing facility claim |
| Hospice Services | Number of eligible beneficiaries with at least one hospice claim |
| Medical Rehabilitation Services | Number of eligible beneficiaries with at least one medical rehabilitation claim |

APPENDIX E: TIME FRAMES AND FORMULAE

| Time Frames | |
|-------------|--------------------------|
| Quarters | Dates |
| Q1 | January 1 to March 31 |
| Q2 | April 1 to June 30 |
| Q3 | July 1 to September 30 |
| Q4 | October 1 to December 31 |

Formulae

$$\text{Incidence} = \frac{\text{(Number of new cases in a time frame, not present in prior year)}}{\text{(Total beneficiaries at risk during the time frame)}}$$

$$\text{Prevalence} = \frac{\text{(Number of cases in a time frame)}}{\text{(Total beneficiaries in the population during the time frame)}}$$

$$\text{Utilization} = \frac{\text{(Number of beneficiaries or measures with specific service utilization)}}{\text{(Total beneficiaries)}}$$

$$\text{Relative change} = \frac{\text{(Current rate-Former rate)}}{\text{(Former rate)}}$$

REFERENCES

1. Weiss MG, Saraceno B, Saxena S, van Ommeren M. Mental health in the aftermath of disasters: consensus and controversy. *The Journal of Nervous and Mental Disease*. 2003 Sep; 191(9):611-615.
2. Foa EB, Stein DJ, McFarlane AC. Symptomatology and psychopathology of mental health problems after disaster. *The Journal of Clinical Psychology* [Internet]. 2006;[cited 16 Sep 2013];67 Suppl 2:15-25.
3. Wang PS, Gruber MJ, Powers RE, Schoenbaum M, Speier AH, Wells KB, Kessler RC. Mental health service use among hurricane Katrina survivors in the eight months after the disaster. *Psychiatry Services* [Internet]. 2007 Nov [cited 16 Sep 2013]; 58(11):1403-1411. Available from: <http://ps.psychiatryonline.org/data/Journals/PSS/3824/07ps1403.pdf>
4. Voelker R. Post-katrina mental health needs prompt group to compile disaster medicine guide. *JAMA*. 2006 Jan [cited 2013 Sep 17]; 295(3):259-260.
5. Centers for Medicare & Medicaid Services. Medicare Claims Database, Parts A and B, January 1, 2011 – March 31, 2013. Baltimore (MD): CMS, Department of Health and Human Services. Accessed: September 15, 2013.
6. U.S. Department of Commerce: United States Census Bureau, American Fact Finder [Internet]. Washington (DC): U.S. Department of Commerce. Median Income in the Past 12 Months (in 2012 Inflation-Adjusted Dollars); 2012 [cited 15 Sep 2013]; [about 2 screens]. Available from: http://factfinder2.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_12_1YR_S1903&prodType=table
7. Buccaneer, A General Dynamics Company. Chronic Condition Data Warehouse: Additions and Access – Task Order 10 New Clinical Conditions: Requirements and Processing [Internet]. [unknown]: Buccaneer, A General Dynamics Company. 2013 May 22 [cited 17 Sep 2013]. Available from: https://www.ccwdata.org/cs/groups/public/documents/document/clin_cond_algo_req_proc.pdf
8. Healthcare Cost and Utilization Project (H-CUP). Clinical Classifications Software (CCS) for ICD-9-CM [Internet]. Rockville (MD): Agency for Healthcare Research and Quality; Nov 2013 [15 Sep 2013]. Available from: <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>
9. Centers for Medicare & Medicaid Services. Mental Health Services Billing Guide, April 2013. Hingham (MA): NHIC, Corp. Apr 2013. 40 p.

| | |
|--|----|
| Executive Summary | |
| Figure 1: Snapshot of Atlantic County | 8 |
| Demographics | |
| Figure 2: Total Medicare FFS Beneficiary Population by County | 9 |
| Figure 3: Percent of Medicare FFS Beneficiary Population by Gender by County | 10 |
| Figure 4: Percent of Medicare FFS Beneficiary Population by Race by County | 11 |
| Figure 5: Percent of Medicare FFS Beneficiary Population by Age by County | 12 |
| Figure 6: 2012 Median Household Income (65 years and above) | 12 |
| Prevalence and Incidence | |
| Figure 7: Prevalence of Selected Behavioral Health Conditions | 14 |
| Figure 8: Prevalence of Depression or Proxy Disorders | 15 |
| Figure 9: Prevalence of Depression or Proxy Disorders in 10 Counties | 16 |
| Figure 10: Atlantic County Prevalence of Depression or Proxy Disorders | 17 |
| Figure 11: Quarterly New Incidence of Depression or Proxy Disorders | 18 |
| Figure 12: Quarterly Relative Change in New Incidence of Depression or Proxy Disorders | 18 |
| Figure 13: Yearly Prevalence of Depression or Proxy Disorders | 19 |
| Figure 14: Yearly Relative Change in Prevalence of Depression or Proxy Disorders | 19 |
| Figure 15: Prevalence of Depression | 20 |
| Figure 16: Quarterly New Incidence of Depression | 21 |
| Figure 17: Quarterly Relative Change in New Incidence of Depression | 21 |
| Figure 18: Yearly Prevalence of Depression | 22 |
| Figure 19: Yearly Relative Change in Prevalence of Depression | 22 |
| Figure 20: Prevalence of Anxiety Disorders | 23 |
| Figure 21: Prevalence of Adjustment Disorders | 23 |
| Figure 22: Prevalence of PTSD | 24 |
| Figure 23: Quarterly New Incidence of PTSD | 25 |
| Figure 24: Quarterly Relative Change in New Incidence of PTSD | 25 |
| Figure 25: Yearly Prevalence of PTSD | 26 |
| Figure 26: Yearly Relative Change in Prevalence of PTSD | 26 |
| Figure 27: Prevalence of Alcohol or Substance Abuse | 27 |
| Figure 28: Quarterly New Incidence of Alcohol or Substance Abuse | 28 |
| Figure 29: Quarterly Relative Change in New Incidence of Alcohol or Substance Abuse | 28 |
| Figure 30: Yearly Prevalence of Alcohol or Substance Abuse | 29 |
| Figure 31: Yearly Relative Change in Prevalence of Alcohol or Substance Abuse | 29 |
| Figure 32: Prevalence of Substance Abuse | 30 |
| Figure 33: Quarterly New Incidence of Substance Abuse | 31 |
| Figure 34: Quarterly Relative Change in New Incidence of Substance Abuse | 31 |
| Figure 35: Yearly Prevalence of Substance Abuse | 32 |
| Figure 36: Yearly Relative Change in Prevalence of Substance Abuse | 32 |
| Figure 37: Rate of Suicide and Intentional Self-Inflicted Injury | 33 |

| | |
|--|----|
| Figure 38: Quarterly New Incidence of Suicide and Intentional Self-Inflicted Injury | 34 |
| Figure 39: Quarterly Relative Change in New Incidence of Suicide and Intentional Self-Inflicted Injury | 34 |
| Figure 40: Yearly Rate of Suicide and Intentional Self-Inflicted Injury | 35 |
| Figure 41: Yearly Relative Change in Rate of Suicide and Intentional Self-Inflicted Injury | 35 |
| Risk Factors for Depression or Proxy Disorders | |
| Figure 42: Prevalence of Top Five Risk Factors for Depression or Proxy Disorders | 36 |
| Figure 43: Prevalence of Top Five Risk Factors for Depression or Proxy Disorders in 10 Counties | 37 |
| Figure 44: Atlantic County Prevalence of Top Five Risk Factors for Depression or Proxy Disorders | 38 |
| Figure 45: Prevalence of Alzheimer's Disease and Related Disorders or Senile Dementia | 39 |
| Figure 46: Prevalence of Sleep Disturbance | 39 |
| Figure 47: Prevalence of Substance or Alcohol Abuse, or Tobacco Use | 40 |
| Figure 48: Prevalence of Hip/Pelvic Fractures | 40 |
| Figure 49: Prevalence of Amputations | 41 |
| Outpatient Behavioral Health Services | |
| <i>Assessments</i> | |
| Figure 50: Utilization of Outpatient Behavioral Health Services– Assessments | 43 |
| Figure 51: Utilization of Outpatient Behavioral Health Services – Therapies | 44 |
| Figure 52: Depression Screening | 45 |
| Figure 53: Depression Screening in 10 Counties | 46 |
| Figure 54: Atlantic County Depression Screening | 47 |
| Figure 55: Quarterly Depression Screening | 48 |
| Figure 56: Diagnostic Psychological Tests | 49 |
| Figure 57: Health and Behavior Assessment/Intervention | 50 |
| Figure 58: Neuropsychological Tests | 51 |
| Figure 59: Quarterly Neuropsychological Tests | 52 |
| Figure 60: Psychiatric Diagnostic Procedures | 53 |
| Figure 61: Quarterly Psychiatric Diagnostic Procedures | 54 |
| <i>Therapies</i> | |
| Figure 62: Individual Psychotherapy | 55 |
| Figure 63: Quarterly Individual Psychotherapy | 56 |
| Figure 64: Family Psychotherapy | 57 |
| Figure 65: Group Psychotherapy | 58 |
| Figure 66: Electronconvulsive Therapy | 59 |
| Figure 67: Biofeedback Therapy | 60 |

Inpatient Services

Admissions

| | |
|---|----|
| Figure 68: Utilization of Inpatient Health Services | 61 |
| Figure 69: Utilization of Inpatient Health Services Within 30 Days of Discharge | 62 |
| Figure 70: Psychiatric Hospital Admissions | 63 |
| Figure 71: Yearly Psychiatric Hospital Admissions | 63 |
| Figure 72: Acute Care Hospital Admissions | 64 |
| Figure 73: Yearly Acute Care Hospital Admissions | 64 |
| Figure 74: Observation Stays | 65 |
| Figure 75: Yearly Observation Stays | 65 |
| Figure 76: Emergency Department Visits | 66 |
| Figure 77: Yearly Emergency Department Visits | 66 |
| Figure 78: 30-Day Hospital Readmissions | 67 |
| Figure 79: Yearly 30-Day Hospital Readmissions | 67 |
| Figure 80: Observation Stays Within 30 Days of Discharge | 68 |
| Figure 81: Yearly Observation Stays Within 30 Days of Discharge | 68 |
| Figure 82: Emergency Department Visits Within 30 Days of Discharge | 69 |
| Figure 83: Yearly Emergency Department Visits Within 30 Days of Discharge | 69 |

Other Settings

| | |
|--|----|
| Figure 84: Utilization of Health Services | 70 |
| Figure 85: Utilization of Home Health Agency Services | 71 |
| Figure 86: Yearly Utilization of Home Health Agency Services | 71 |
| Figure 87: Utilization of Skilled Nursing Facility Services | 72 |
| Figure 88: Yearly Utilization of Skilled Nursing Facility Services | 72 |
| Figure 89: Utilization of Hospice Services | 73 |
| Figure 90: Yearly Utilization of Hospice Services | 73 |
| Figure 91: Utilization of Medical Rehabilitation Services | 74 |
| Figure 92: Yearly Utilization of Medical Rehabilitation Services | 74 |

INDEX OF FIGURES

This page left intentionally blank



557 Cranbury Road, Suite 21
East Brunswick, NJ 08816
T 732-238-5570 • F 732-238-7766
www.hqsi.org