


# Enhancing Coordination of Behavioral Health Services after Superstorm Sandy: Planning for Future Disasters

---

## Final Data Profile: Toms River Community Medicare Fee-for-Service Beneficiaries

*Demographics, Behavioral Health Conditions, and Utilization of Health Services (Medicare Fee-for-Service Beneficiaries)*

July 9, 2014



This Page Left Intentionally Blank

<b>Preface</b> .....	1
<b>Introduction</b> .....	2
<b>HQSI Project Team</b> .....	7
<b>Acknowledgements</b> .....	8
<b>Executive Summary</b> .....	9
<b>Demographics</b> .....	11
• Total Medicare FFS Beneficiaries .....	11
• Percent of Medicare FFS Beneficiaries in the General Population in 2012 .....	11
• Percent of Medicare FFS Beneficiary Population by Female .....	11
• Percent of Medicare FFS Beneficiary Population by Race .....	12
• Percent of Medicare FFS Beneficiary Population by Age.....	12
• Median Household Income (65 Years and Above) .....	12
<b>Behavioral Health Conditions</b> .....	13
• Prevalence and Incidence.....	13
– Summary .....	13
– Depression or Proxy Disorders .....	16
– Depression.....	22
– Anxiety Disorders .....	23
– Adjustment Disorders .....	24
– Post-Traumatic Stress Disorder.....	25
– Alcohol or Substance Abuse .....	26
– Substance Abuse Alone .....	27
– Suicide and Intentional Self-Inflicted Injury .....	28
• Risk Factors for Depression or Proxy Disorders.....	29
– Summary .....	29
– Any of the Top Five Risk Factors for Depression or Proxy Disorders .....	30
– Alzheimer's Disease and Related Disorders or Senile Dementia.....	33
– Sleep Disturbance.....	33
– Substance or Alcohol Abuse or Tobacco Use .....	33
– Hip/Pelvic Fractures .....	34
– Amputations .....	34
<b>Utilization</b> .....	35
• Utilization of Outpatient Behavioral Health Services .....	35
– Assessments .....	35
› Summary .....	35
› Depression Screening .....	35

# TABLE OF CONTENTS

---

› Neuropsychological Tests .....	40
› Psychiatric Diagnostic Procedures .....	42
– Therapies .....	44
› Summary .....	44
› Individual Psychotherapy .....	45
› Family Psychotherapy .....	47
› Group Psychotherapy .....	48
• Inpatient Health Services .....	49
– Summary .....	49
– Psychiatric Hospital Admissions .....	50
– Acute Care Hospitals .....	51
› Admissions .....	51
› Observation Stays .....	52
› Emergency Department Visits .....	53
– Within 30 Days of Acute Care Hospital Discharge .....	54
› Summary .....	54
› 30-Day Hospital Readmissions .....	55
› Observation Stays Within 30 Days of Discharge .....	56
› Emergency Department Visits Within 30 Days of Discharge .....	57
– Other Settings .....	58
› Summary .....	58
› Home Health Agency Services .....	59
› Skilled Nursing Facility Services .....	60
› Hospice Services .....	61
› Medical Rehabilitation Services .....	62
• Listing of Major Health Providers .....	63
• Toms River Community Providers .....	64
• Appendix A: Behavioral Health Conditions .....	65
• Appendix B: Risk Factors for Depression or Proxy Disorders .....	68
• Appendix C: Utilization of Outpatient Mental Health Services .....	72
• Appendix D: Utilization of Services – Inpatient and Other Settings .....	74
• Appendix E: Time Frames and Formulae .....	76
• Appendix F: References .....	77
• Appendix G: Provider Summary Tables and Provider Listings .....	78
<b>Index of Figures .....</b>	<b>81</b>

**O**n October 29, 2012, Superstorm Sandy hit the Eastern Seaboard, impacting more than a dozen states. New Jersey, which took the brunt of the storm along its densely populated coastline, was devastated. Thousands of residents were displaced, their homes and communities damaged or destroyed.

Lessons learned from prior natural disasters showed that victims of storms like Superstorm Sandy are often at an elevated risk for acute or behavioral health issues such as post-traumatic stress disorder (PTSD), depression, and substance abuse.<sup>1,2</sup> While disaster-related issues subside over time, evidence shows that victims can experience a prolonged period of elevated risk, especially those with pre-existing mental health issues.<sup>3</sup> Older adults and disabled residents with mental health conditions are at increased risk of deteriorating health, depression, increased isolation, and breakdown in the continuum of health care. Additionally, past natural disasters also show that access to informational resources on disaster-related mental health disorders, outcomes, and service utilization are important factors to consider.<sup>4,5</sup>

This final community profile can help healthcare professionals learn more about the behavioral health status and utilization patterns of Medicare Fee-for-Service (FFS) beneficiaries before and after Superstorm Sandy. As such, it may be a useful tool in planning for future disasters. The profile explores community-level health status and health determinants of post-disaster spikes in behavioral health issues and treatments. This last update includes one more quarter of comprehensive post- Sandy data than the previous profile, which was published in May 2014.

*Enhancing Coordination of Behavioral Health Services after Superstorm Sandy: Planning for Future Disasters* is a Special Innovation Project funded by the Centers for Medicare & Medicaid Services (CMS). As part of this project, Healthcare Quality Strategies, Inc. (HQSI), the CMS quality improvement organization (QIO) for New Jersey, studied data on prevalence and incidence of selected behavioral health conditions, the utilization of health services, and demographic information from the Medicare claims database for Medicare FFS beneficiaries residing in the 10 New Jersey FEMA-declared disaster counties after Superstorm Sandy. From its analysis, HQSI created data profiles for 10 FEMA-designated counties, as well as a subset of 10 selected communities.

This final profile is the last update planned for the Toms River community and includes data from January 1, 2011 to December 31, 2013. The Toms River community was selected because it had high rates of Medicare FFS beneficiaries both with and at risk for depression or proxy disorders prior to Superstorm Sandy. This profile can be used to determine and compare the community's prevalence and incidence of the selected behavioral health conditions and utilization of services to its county rate before and after Superstorm Sandy.

The data in the profile is based on Medicare FFS claims data and provides a glimpse into the prevalence and incidence of selected behavioral health conditions and risk factors for depression, as well as the utilization of Medicare-covered behavioral health services among Medicare beneficiaries residing in the community before and after Superstorm Sandy. Since patients with behavioral health conditions may receive other health services because of medical problems caused by their behavioral health conditions or they may avoid utilizing behavioral health services, this report also looks at the utilization of non-behavioral health services.

The county and community profiles are being shared with state and local governments and agencies, health care providers, community-based organizations, and the research community to support a community-based approach to enhance the coordination of behavioral health services after a natural disaster, and to increase utilization of the Medicare depression screening benefit which became a covered service in October 2011. This benefit is important for victims of major disasters like Superstorm Sandy who are often at an elevated risk for behavioral health issues and can experience a prolonged period of elevated risk after a disaster. Older adults and disabled residents with behavioral health conditions in particular are at increased risk of deteriorating health, depression, increased isolation, and breakdown in the continuum of health care. They are also less likely to report symptoms, which a depression screening can capture.

## WHAT'S NEW IN THIS UPDATE

This second updated profile shows four quarters of post-Sandy data, with the most updated claims from January 2011 to December 2013. The profile compares the 12-month rates from the year before and after the storm. In this profile, we define October 2011 to September 2012 as the year before Superstorm Sandy, October 1, 2012 through December 31, 2012 as the time frame when Superstorm Sandy occurred, and January 2013 to December 2013 as the year after the storm.

## HOW TO USE THIS PROFILE

This profile includes an analysis of the eight behavioral health conditions which, based on literature review and feedback from the subject matter experts consulted for this project, were found to increase after natural disasters.

This profile is divided into the following sections, each of which is preceded by a user-friendly overview:

- Demographics (page 11)
- Prevalence and incidence of behavioral health conditions (page 13)
- Risk factors for depression or proxy disorders (page 29)
- Utilization of outpatient behavioral health assessments (page 35)
- Utilization of outpatient behavioral health therapies (page 44)
- Utilization of inpatient health services (page 49)
- Utilization of inpatient health services within 30 days of acute care hospital discharge (page 54)
- Utilization of other settings (page 58)
- Listing of major health providers (page 63)

Here are some additional tips for using this profile:

- Use the Executive Summary (pages 9-10) for a quick overview of this profile's key points, as well as a snapshot table that summarizes the prevalence of the selected behavioral health conditions and utilization of behavioral health services before and after Sandy
- Use the Behavioral Health Conditions section (pages 13-34) for in-depth analyses and graphical comparison on the prevalence and incidence of eight behavioral health conditions before and after Superstorm Sandy
- Use the New Jersey and county maps to: identify areas with higher rates of Medicare FFS beneficiaries at risk for depression and proxy disorders (pages 31-32); and areas with low utilization of the depression screening benefit (pages 38-39)

## METHODOLOGY

Each community profile compares the community's statistics to the statistics of the county where each community is located. Primary data sources include Medicare FFS Part A and Part B claims, the Medicare enrollment database, and U.S. Census data. The Medicare enrollment database includes basic demographic statistics such as age, gender, and race while the U.S. Census data provides a proxy indicator (average household income) for socio-economic status.<sup>6</sup> Based on the ICD-9-CM (International Classification of Disease, Ninth Revision, Clinical Modification), CPT (Current Procedural Terminology), or HCPCS (Healthcare Common Procedure Coding System) codes in Medicare Part A and Part B claims, beneficiaries were identified for diseases/conditions related to behavioral health conditions such as depression. Appendices A through G contain documentation, technical notes, codes, algorithms, data sources, and references.

Medicare Part A claims were also used to analyze utilization of health services in acute care hospitals, skilled nursing facilities, medical rehabilitation facilities, home health agencies, hospice, and inpatient psychiatric facilities. Medicare Part A and Part B claims provide information on the utilization of mental health outpatient services for assessment (e.g., depression screening, diagnostic psychological tests) and treatment (e.g., individual psychotherapy, biofeedback therapy).

To identify beneficiaries with an elevated risk of depression after the storm, HQSI conducted a literature review of risk factors for depression (see Appendix B). Previous studies identified psychosocial and biological factors, increased age, history of cancer, Parkinson's disease, Alzheimer's disease, changes in mental function, and medication side effects as risk factors for developing depression. Based on findings from the literature review and factors available through Medicare claims, logistic regression analysis was conducted with Medicare claims, and the top five risk factors (Alzheimer's disease and related disorders or senile dementia, hip/pelvic fractures, amputations, substance or alcohol abuse or tobacco use, and sleep disturbance) were used to identify beneficiaries with high risk for developing depression or proxy disorders (i.e., anxiety and adjustment disorders).



## MEASUREMENT TIME FRAMES

This profile includes data from January 1, 2011 through December 31, 2013. Results are presented using different charts and measurement time frames as follows:

- Annual bar charts show the annual rates in the year before (October 1, 2011 to September 30, 2012) and after (January 1, 2011 to December 31, 2013) Superstorm Sandy. Statistics on demographics, prevalence of behavioral health conditions, and utilization of health services are presented for this 12-month period. These statistics allow for comparison before and after Superstorm Sandy
- Annual trend charts with rolling quarters for the behavioral health conditions and utilization statistics are included to adjust for seasonal variation and to examine possible changes in the year before and after Superstorm Sandy. The time period includes nine data points from January 1, 2011 to December 31, 2013
- Annual percent change (relative change) bar charts show relative increase or decrease in rates from the year before and after Superstorm Sandy. These statistics allow for comparison of the community to the county and to analyze the potential impact of Superstorm Sandy
- Quarterly new incidence charts for eight behavioral health conditions include eight quarters of data from January 1, 2012 to December 31, 2013. This allows for the identification of new cases in a given quarter when compared to the prior year
- Quarterly line charts show the trend in the utilization of depression screening for eight quarters from January 1, 2012 to December 31, 2013

### DATA CONSIDERATIONS

There are now four quarters of post-storm data available, which are reflected in this final update. The claims data processing lag of at least six months, coupled with the one-year project time frame, reduces the optimal time frame for more accurate estimation of post-Sandy effects.

Identification of beneficiaries with behavioral health conditions is based on diagnoses being reported in Medicare FFS claims and could result in underestimation. There is no accurate way to identify when certain health conditions began and ended when claims data is used.

According to the subject matter experts consulted for this project, unlike other conditions, behavioral health issues are often underdiagnosed in our society and the stigma associated with behavioral health conditions may prevent people from seeking care in mental health facilities. The subject matter experts also indicated that estimating the prevalence of depression using claims data can be particularly difficult as depression is often undiagnosed or not documented. Depression can be present with symptoms of anxiety and adjustment disorders. Based on this feedback, a combination measure named “depression or proxy disorders” was created to estimate prevalence and incidence of depression. If a patient had at least one of the three conditions reported in Medicare claims, he/she was flagged as having depression or proxy disorders.

This community profile can be used to compare the prevalence and incidence rates of eight selected behavioral health conditions (see page 19) based on the ICD-9-CM codes through the analysis of Medicare claims. This profile may be used to prioritize and plan community and county preparation for the care, tracking, and monitoring of Medicare beneficiary behavioral health status and health care utilization patterns.

This is the final update of these data profiles which includes one more quarter of data than the previous profile during the post-Superstorm Sandy time period as the project ends on July 31, 2014.

**Martin P. Margolies**  
Chief Executive Officer

**Mary Jane Brubaker, MCIS**  
Chief Operating Officer

**Diane Babuin, MS, CPHQ**  
Director, Quality Improvement and Communications

**Ya-ping Su, PhD**  
Director, Research and Analysis

**Suzanne Dalton, RN, BS, EdM**  
Project Manager

**Andrew Miller, MD, MPH**  
Medical Director

---

**Mona Abdalla, BA**  
Administrative Associate

**Christine Aisenberg, BA**  
Proofreader

**Kathy Brown, BS**  
InDesign Specialist

**Zhengyu Bu, MS**  
Health Services Research Analyst

**Sue Chen, MS**  
Statistician

**Wei-Yi Chung, MS**  
Database Administrator

**Barbara Coleman**  
Administrative Associate

**Dawn Cullen, BA**  
Communications Specialist

**Ashley Dopp, BA**  
Communications Specialist

**Karen Hale, MEd**  
Community Liaison

**Kim Karnell, BS**  
Information Specialist

**Janet Knoth, BS, RN, CHPN, CPHQ**  
Quality Improvement Specialist

**Judy Miller, MS, RN**  
Quality Improvement Specialist

**Olubukunola Oyedele, MPH**  
Community Liaison

**Rita Pascale**  
Administrative Associate

**Barbara Perzyna, BS**  
Visual Communications Specialist

**Ziphora Sam, MPH**  
Epidemiologist

**Marianne Sagarese, BSN, RN**  
Quality Improvement Specialist

**Nicole Skyer-Brandwene, MS, RPh, BCPS**  
Quality Improvement Specialist

# ACKNOWLEDGEMENTS

---

Special thanks to the subject matter experts who assisted with the project by providing feedback and guidance to the HQSI project team.

**Carol Benevy, MSW**

New Jersey Hope and Healing Project  
Barnabas Health Institute for Prevention

**Mary Ditri, MA, CHCC**

New Jersey Hospital Association

**Adrienne Fessler-Belli, MSW, LCSW**

New Jersey Department of Human Services  
Disaster & Terrorism Branch

**Mark Firth, MA, MSW**

New Jersey Department of Human Services  
Division of Mental Health and Addiction Services

**Mary Goepfert, MPA, APR, CPM**

New Jersey Group for Access and Integration Needs in Emergencies and Disasters

**Sheldon Green**

New Jersey Primary Care Association

**Connie Greene, MA, CAS, CSW, CPS**

Barnabas Health Institute for Prevention

**Bob Kley**

Mental Health Association in New Jersey, Inc.

**Lynn Kovitch, MEd**

New Jersey Department of Human Services  
Division of Mental Health and Addiction Services

**Karen McCoy, RN, BSN**

Home Care Association of New Jersey

**Elyse Perweiler, MPP, RN**

NJ Institute for Successful Aging

**Lynn Stefanowicz, MA, LCSW**

Meridian Behavioral Health

**Megan Sullivan, LPC, LCADC, DRCC**

New Jersey Department of Human Services  
Disaster & Terrorism Branch

**Pete Summers**

The New Jersey Association of County and City Health Officials (NJACCHO)

## KEY OBSERVATIONS

The following observations show the Toms River community's percent change and rates after Superstorm Sandy among Medicare FFS beneficiaries.

1. The Toms River community experienced a larger decrease in adjustment disorders (14.82%) than Ocean County. There was a relative increase in depression or proxy disorders (4.13%), anxiety disorders (12.99%), alcohol or substance abuse (6.13%), and PTSD (9.69%) in the community.
2. The demographic breakdown of beneficiaries residing in the Toms River community, who experienced higher rates of depression or proxy disorders, is as follows: Hispanic (333.33 per 1,000 beneficiaries), females (258.52 per 1,000 beneficiaries), and those below age 65 (377.42 per 1,000 beneficiaries).
3. The Toms River community experienced a larger decrease in any of the top five risk factors for depression or proxy disorders (10.50%) than Ocean County. This also includes a larger decrease in substance or alcohol abuse or tobacco use (10.29%), Alzheimer's disease and related disorders or senile dementia (22.51%), hip/pelvic fractures (16.59%), and amputations (49.52%).
4. Annual depression screening in the Toms River community increased from 8.78 per 1,000 beneficiaries before the storm to 13.51 per 1,000 after the storm.
5. The Toms River community experienced a larger decrease in psychiatric diagnostic procedures (25.64%) than Ocean County.
6. The Toms River community experienced a larger decrease in the utilization of family psychotherapy (33.19%) than Ocean County.
7. The Toms River community experienced a larger increase in observation stays (520.97%) and observation stays that occurred within 30 days of discharge (39.96%) than Ocean County.
8. The Toms River community experienced a larger decrease in the utilization of skilled nursing facility services (12.07%) than Ocean County.

# EXECUTIVE SUMMARY


The *Snapshot of the Toms River Community* (Figure 1) summarizes the prevalence of the behavioral health conditions as well as risk factors for depression or proxy disorders analyzed for this profile. This *Snapshot* also lists the most frequently performed behavioral health assessments and therapies in the Toms River community compared to the average of Ocean County. The non-behavioral health utilization measures that were calculated for this profile are not included in the *Snapshot*.

**Figure 1. Snapshot of Toms River**


Behavioral Health Disorders	Prevalence per 1,000 Medicare FFS Beneficiaries					
	Toms River			Ocean County		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Depression or Proxy Disorders	212.30	221.07	4.13	208.85	220.98	5.81
• Depression alone	134.86	134.83	-0.02	131.66	135.96	3.27
• Anxiety Disorders alone	129.13	145.90	12.99	125.55	142.68	13.64
• Adjustment Disorders alone	27.26	23.22	-14.82	28.88	26.27	-9.04
Alcohol or Substance Abuse	37.68	39.99	6.13	35.33	42.09	19.13
• Substance Abuse alone	24.59	24.54	-0.20	20.67	22.36	8.18
PTSD	6.19	6.79	9.69	5.86	6.70	14.33
Suicide and Intentional Self-Inflicted Injury	6.49	5.96	-8.17	5.33	5.34	0.19
Any of the Top Five Risk Factors* for Depression or Proxy Disorders						
• Any of the Top Five Risk Factors	170.78	152.85	-10.50	161.01	156.28	-2.94
• Substance or Alcohol Abuse or Tobacco Use	111.70	100.21	-10.29	104.83	102.85	-1.89
• Alzheimer’s Disease and related disorders or Senile Dementia	46.38	35.94	-22.51	39.75	33.91	-14.69
• Sleep Disturbance	24.27	24.35	0.33	25.76	27.77	7.80
• Hip/Pelvic Fractures	8.50	7.09	-16.59	7.72	6.91	-10.49
• Amputations	1.05	0.53	-49.52	1.02	0.74	-27.45
Behavioral Health Services	Utilization per 1,000 Medicare FFS Beneficiaries					
	Toms River			Ocean County		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Assessments						
• Depression Screening**	8.78	13.51	53.87	9.50	16.27	71.26
• Psychiatric Diagnostic Procedures	57.06	42.43	-25.64	54.39	43.28	-20.43
• Neuropsychological Tests	12.31	11.13	-9.59	9.77	12.54	28.35
Therapy						
• Individual Psychotherapy	46.41	49.05	5.69	48.83	52.21	6.92
• Family Psychotherapy	2.32	1.55	-33.19	2.70	2.20	-18.52
• Group Psychotherapy	1.28	1.05	-17.97	1.75	1.43	-18.29
Psychiatric Hospital Admissions	7.46	7.81	4.69	7.67	6.97	-9.13

\*The top five risk factors were identified based on findings from a literature review (Appendix B) and factors available through Medicare claims. Logistic regression analysis was conducted with Medicare claims.

\*\*Depression Screening comparison time frames are different (January 1, 2012 – December 31, 2012 vs. January 1, 2013 – December 31, 2013) due to availability of depression screening data starting in January 2012.

Medicare FFS Demographics		Toms River	
<b>At A Glance</b> (January 1, 2013 – December 31, 2013)			
Total Medicare FFS Population	32,206		
Females	18,539 (57.56%)		
Males	13,667 (42.44%)		
White	30,739 (95.44%)		
Black	478 (1.48%)		
Asian	257 (0.80%)		
Hispanic	190 (0.59%)		
Other	542 (1.68%)		
Average Age	73.94		
ZIP Codes	08735, 08751, 08753, 08754, 08755, 08756, 08757		
Source: Medicare Claims Database			


**FIGURE 2. TOTAL MEDICARE FFS BENEFICIARIES\***



\* Total beneficiaries who were under Medicare FFS coverage for at least one month during the time frame.

The total Medicare FFS population of the Toms River community prior to Superstorm Sandy was 34,287. After the storm, this number decreased to 32,206.

**FIGURE 3. PERCENT OF MEDICARE FFS BENEFICIARIES IN THE GENERAL POPULATION IN 2012\***



\* Source: Medicare denominator file CY 2012, U.S. Census Bureau, American Cancer Survey (ACS), 2012 <http://www.census.gov/>.

Medicare FFS beneficiaries made up 25.03% of the Toms River community in calendar year 2012.

**FIGURE 4. PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY FEMALE**

	10/1/11 – 9/30/12	1/1/13 – 12/31/13	Absolute Change*
Toms River	57.91	57.56	-0.35
Ocean County	57.16	56.96	-0.20

\* Due to rounding, the absolute change may not be the same as the difference subtracted from the two time frames shown.

Prior to the storm, females made up 57.91% of the entire Medicare FFS population residing in the Toms River community and males made up 42.09%. After the storm, the female beneficiary population decreased to 57.56% and males increased to 42.44%.

**FIGURE 5. PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY RACE**

Race	Toms River			Ocean		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	Absolute Change*	10/1/11 – 9/30/12	1/1/13 – 12/31/13	Absolute Change*
White	95.76	95.44	-0.31	95.60	95.30	-0.30
Black	1.54	1.48	-0.06	1.96	1.92	-0.04
Hispanic	0.56	0.59	0.03	0.57	0.56	-0.01
Asian	0.76	0.80	0.03	0.57	0.58	0.01
Other	1.38	1.68	0.31	1.30	1.65	0.34

\* Due to rounding, the absolute change may not be the same as the difference subtracted from the two time frames shown.

Both before and after Superstorm Sandy, the majority of Medicare FFS beneficiaries residing in the Toms River community were White followed by Black, Asian, and Hispanic.

**FIGURE 6. PERCENT OF MEDICARE FFS BENEFICIARY POPULATION BY AGE\***


Age	Toms River			Ocean		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	Absolute Change**	10/1/11 – 9/30/12	1/1/13 – 12/31/13	Absolute Change**
<65	11.44	11.12	-0.33	11.44	11.28	-0.17
65 – 74	39.06	41.12	2.05	43.25	44.44	1.19
75 – 84	30.13	29.01	-1.13	29.02	28.50	-0.52
85 and Above	19.36	18.76	-0.60	16.28	15.78	-0.50
Average Age	74.10	73.94	-0.17	73.34	73.24	-0.11

\* Age calculated as end date of time frame or date of death minus birth date.

\*\* Due to rounding, the absolute change may not be the same as the difference subtracted from the two time frames shown.

Both before and after Superstorm Sandy, the largest age group of Medicare FFS beneficiaries residing in the Toms River community was between ages 65 and 74 years old, followed by beneficiaries between 75 and 84 years old. The average age of beneficiaries residing in this community decreased from 74.10 years old prior to the storm to 73.94 years old after the storm.

**FIGURE 7. 2012 MEDIAN HOUSEHOLD INCOME (65 YEARS AND ABOVE)**



According to U.S. Census Data from 2012, residents aged 65 and older residing in the Toms River community had a median household income of \$43,250. This was higher than the average income among seniors residing in all of Ocean County.

Source: U.S. Census Bureau, American Community Survey (ACS), 2012 <http://www.census.gov/>.



## PREVALENCE AND INCIDENCE

Using Medicare FFS claims data, eight behavioral health conditions were analyzed: depression or proxy disorders, depression, adjustment disorder, anxiety disorder, post-traumatic stress disorder (PTSD), alcohol or substance abuse, substance abuse alone, and suicide and intentional self-inflicted injury. These conditions were chosen based on literature review and feedback from subject matter experts.

Claims data can underestimate the real prevalence and incidence of depression in the population and individuals with depression could be diagnosed as having anxiety or adjustment disorders, as noted by the subject matter experts consulted for this project. Therefore, HQSI created a combination measure for depression (depression or proxy disorders) which includes beneficiaries who were reported for either depression, anxiety, or adjustment disorders.

The behavioral health data from January 1, 2011 to December 31, 2013 for these different measures were calculated to quantify condition occurrence:

1. The annual prevalence bar chart compares rates in two annual time frames
2. New incidence in a quarter for the specified condition that was not present in the prior 12 months (Q1 2012 – Q4 2013)
3. The yearly prevalence of the condition with quarterly rolling trends to account for seasonal variation


Refer to Appendix A for measurement calculation and Appendix E for quarterly time frames and formulae.

### Summary

<b>Figure 8. Percent Change of Prevalence of Selected Behavioral Health Conditions per 1,000 Medicare FFS Beneficiaries</b>						
	<b>Toms River</b>			<b>Ocean</b>		
	<b>10/1/11 – 9/30/12</b>	<b>1/1/13 – 12/31/13</b>	<b>% Change</b>	<b>10/1/11 – 9/30/12</b>	<b>1/1/13 – 12/31/13</b>	<b>% Change</b>
Depression or Proxy Disorders	212.30	221.07	4.13	208.85	220.98	5.81
• Depression	134.86	134.83	-0.02	131.66	135.96	3.27
• Anxiety	129.13	145.90	12.99	125.55	142.68	13.64
• Adjustment	27.26	23.22	-14.82	28.88	26.27	-9.04
Alcohol or Substance Abuse	37.68	39.99	6.13	35.33	42.09	19.13
• Substance abuse alone	24.59	24.54	-0.20	20.67	22.36	8.18
PTSD	6.19	6.79	9.69	5.86	6.70	14.33
Suicide and intentional self-inflicted injuries	6.49	5.96	-8.17	5.33	5.34	0.19


The Toms River community experienced a larger decrease in adjustment disorders than Ocean County.

**FIGURE 9. QUARTERLY NEW INCIDENCE TREND OF SELECTED BEHAVIORAL HEALTH CONDITIONS: DEPRESSION OR PROXY DISORDERS\* PER 1,000 MEDICARE FFS BENEFICIARIES**



\* Quarterly new incidence of conditions that were not diagnosed in the prior year.


**FIGURE 10. QUARTERLY NEW INCIDENCE TREND OF OTHER SELECTED BEHAVIORAL HEALTH CONDITIONS\* PER 1,000 MEDICARE FFS BENEFICIARIES**




\* Quarterly new incidence of conditions that were not diagnosed in the prior year.

The charts above reflect quarterly trending in new incidence of the eight selected behavioral health conditions among Medicare FFS beneficiaries residing in the Toms River community.

**FIGURE 11. ANNUAL PREVALENCE TREND OF SELECTED BEHAVIORAL HEALTH CONDITIONS: DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES**



**FIGURE 12. ANNUAL PREVALENCE TREND OF OTHER SELECTED BEHAVIORAL HEALTH CONDITIONS PER 1,000 MEDICARE FFS BENEFICIARIES**



The charts above reflect annual trending in the prevalence of the eight selected behavioral health conditions among Medicare FFS beneficiaries residing in the Toms River community.

## Depression or Proxy Disorders

Figure 13. Demographics of Depression or Proxy Disorders among Medicare FFS Beneficiaries				
	10/1/11 – 9/30/12		1/1/13 – 12/31/13	
	Number of Beneficiaries	Percent (%)	Number of Beneficiaries	Percent (%)
<b>Race</b>				
• White	6,288	96.46	6,471	96.42
• Black	106	1.63	86	1.28
• Hispanic	42	0.64	59	0.88
• Asian	24	0.37	22	0.33
• Other	59	0.91	73	1.09
<b>Gender</b>				
• Males	2,029	31.12	2,180	32.48
• Females	4,490	68.88	4,531	67.52
<b>Age</b>				
• Below 65	1,313	20.14	1,247	18.58
• 65-74	1,984	30.43	2,169	32.32
• 75-84	1,711	26.25	1,747	26.03
• 85 and Above	1,511	23.18	1,548	23.07
<b>Total</b>	<b>6,519</b>	<b>100.00</b>	<b>6,711</b>	<b>100.00</b>


This table displays the number and percentage of Medicare FFS beneficiaries of each race, gender, and age diagnosed with depression or proxy disorders before and after Superstorm Sandy. There were 6,519 beneficiaries residing in the Toms River community diagnosed with depression or proxy disorders before the storm. This increased to 6,711 beneficiaries after the storm.

<b>Figure 14. Depression or Proxy Disorders Rate per 1,000 Medicare FFS Beneficiaries by Demographic Group</b>						
	<b>10/1/11 – 9/30/12</b>			<b>1/1/13 – 12/31/13</b>		
	<b>Numerator</b>	<b>Denominator*</b>	<b>Rate per 1,000 Beneficiaries</b>	<b>Numerator</b>	<b>Denominator*</b>	<b>Rate per 1,000 Beneficiaries</b>
<b>Race</b>						
• White	6,288	29,471	213.36	6,471	29,010	223.06
• Black	106	447	237.14	86	441	195.01
• Hispanic	42	160	262.50	59	177	333.33
• Asian	24	231	103.90	22	239	92.05
• Other	59	401	147.13	73	493	148.07
<b>Gender</b>						
• Males	2,029	12,837	158.06	2,180	12,831	169.90
• Females	4,490	17,872	251.23	4,531	17,527	258.52
<b>Age</b>						
• Below 65	1,313	3,291	398.97	1,247	3,304	377.42
• 65-74	1,984	11,911	166.57	2,169	12,355	175.56
• 75-84	1,711	9,671	176.92	1,747	9,060	192.83
• 85 and Above	1,511	5,837	258.87	1,548	5,640	274.47
<b>Total</b>	<b>6,519</b>	<b>30,710</b>	<b>212.28</b>	<b>6,711</b>	<b>30,359</b>	<b>221.05</b>

\* Total eligible beneficiaries (denominator) computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame


This table displays the rate of Medicare FFS beneficiaries per 1,000 diagnosed with depression or proxy disorders by race, gender, and age both before and after Superstorm Sandy by different demographic groups. The numerator is the number of beneficiaries with a claim for depression or proxy disorders; the denominator is the total number of beneficiaries residing in the community for each group.

**FIGURE 15. DEPRESSION OR PROXY DISORDERS RATE BY RACE PER 1,000 MEDICARE FFS BENEFICIARIES**




Hispanic Medicare FFS beneficiaries have the highest rate of depression or proxy disorders. In the 12 months prior to Superstorm Sandy, 262.50 per 1,000 Hispanic beneficiaries were diagnosed with depression or proxy disorders. After the storm, this rate increased to 333.33 per 1,000 beneficiaries.

**FIGURE 16. DEPRESSION OR PROXY DISORDERS RATE BY GENDER PER 1,000 MEDICARE FFS BENEFICIARIES**




Female Medicare FFS beneficiaries have a higher rate of depression or proxy disorders. In the 12 months prior to Superstorm Sandy, 251.23 per 1,000 female beneficiaries were diagnosed with depression or proxy disorders. After the storm, this rate increased to 258.52 per 1,000 beneficiaries.

**FIGURE 17. DEPRESSION OR PROXY DISORDERS RATE BY AGE GROUP PER 1,000 MEDICARE FFS BENEFICIARIES**




Medicare FFS beneficiaries below the age of 65 have the highest rate of depression or proxy disorders. In the 12 months prior to Superstorm Sandy, 398.97 per 1,000 beneficiaries below the age of 65 were diagnosed with depression or proxy disorders. After the storm, this rate decreased to 377.42 per 1,000 beneficiaries.

**FIGURE 18. DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of depression or proxy disorders among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 212.30 per 1,000 beneficiaries. After the storm, this rate increased to 221.07 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 4.13% relative increase.

**FIGURE 19. QUARTERLY NEW INCIDENCE OF DEPRESSION OR PROXY DISORDERS\* PER 1,000 MEDICARE FFS BENEFICIARIES**



This chart reflects trending of quarterly new incidence of depression or proxy disorders among Medicare FFS beneficiaries residing in the Toms River community.


\* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.

# BEHAVIORAL HEALTH CONDITIONS

**FIGURE 20. PREVALENCE OF DEPRESSION OR PROXY DISORDERS\* PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES**

October 1, 2011 – September 30, 2012

January 1, 2013 – December 31, 2013



The color-coded map of New Jersey depicts prevalence of depression or proxy disorders from high (red) to low (blue) in the 10 FEMA-declared disaster counties before and after Superstorm Sandy.


\* Mapped using ZIP codes of the 10 counties.



**FIGURE 21. OCEAN COUNTY PREVALENCE OF DEPRESSION OR PROXY DISORDERS\* PER 1,000 MEDICARE FFS BENEFICIARIES**

October 1, 2011 – September 30, 2012

January 1, 2013 – December 31, 2013




The color-coded map of Ocean County depicts regional variation of prevalence of depression or proxy disorders from high (red) to low (blue) before and after Superstorm Sandy.

\* Mapped using ZIP codes; may not display all the city names located within the ZIP code.


## Depression

**FIGURE 22. DEPRESSION PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of depression among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 134.86 per 1,000 beneficiaries. After the storm, this rate decreased to 134.83 per 1,000 beneficiaries, reflecting a 0.02% relative decrease.

**FIGURE 23. QUARTERLY NEW INCIDENCE OF DEPRESSION\* PER 1,000 MEDICARE FFS BENEFICIARIES**




This chart reflects trending of quarterly new incidence of depression among Medicare FFS beneficiaries residing in the Toms River community.

\* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


## Anxiety Disorders

**FIGURE 24. ANXIETY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of anxiety disorders among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 129.13 per 1,000 beneficiaries. After the storm, this rate increased to 145.90 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 12.99% relative increase.

**FIGURE 25. QUARTERLY NEW INCIDENCE OF ANXIETY DISORDERS\* PER 1,000 MEDICARE FFS BENEFICIARIES**




This chart reflects trending of quarterly new incidence of anxiety disorders among Medicare FFS beneficiaries residing in the Toms River community.

\* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


## Adjustment Disorders

**FIGURE 26. ADJUSTMENT DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of adjustment disorders among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 27.26 per 1,000 beneficiaries. After the storm, this rate decreased to 23.22 per 1,000 beneficiaries. This change reflects a 14.82% relative decrease, a larger decrease than in Ocean County.

**FIGURE 27. QUARTERLY NEW INCIDENCE OF ADJUSTMENT DISORDERS\* PER 1,000 MEDICARE FFS BENEFICIARIES**




This chart reflects trending of quarterly new incidence of adjustment disorders among Medicare FFS beneficiaries residing in the Toms River community.

\* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


## Post-Traumatic Stress Disorder (PTSD)

**FIGURE 28. PTSD PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of PTSD among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 6.19 per 1,000 beneficiaries. After the storm, this rate increased to 6.79 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 9.69% relative increase.

**FIGURE 29. QUARTERLY NEW INCIDENCE OF PTSD\* PER 1,000 MEDICARE FFS BENEFICIARIES**




This chart reflects trending of quarterly new incidence of PTSD among Medicare FFS beneficiaries residing in the Toms River community.

\* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


## Alcohol or Substance Abuse

**FIGURE 30. ALCOHOL OR SUBSTANCE ABUSE PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of alcohol or substance abuse among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 37.68 per 1,000 beneficiaries. After the storm, this rate increased to 39.99 per 1,000 beneficiaries, reflecting a 6.13% relative increase.

**FIGURE 31. QUARTERLY NEW INCIDENCE OF ALCOHOL OR SUBSTANCE ABUSE\* PER 1,000 MEDICARE FFS BENEFICIARIES**




This chart reflects trending of quarterly new incidence of alcohol or substance abuse among Medicare FFS beneficiaries residing in the Toms River community.

\* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


## Substance Abuse Alone

**FIGURE 32. SUBSTANCE ABUSE ALONE PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of substance abuse alone among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 24.59 per 1,000 beneficiaries. After the storm, this rate decreased to 24.54 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 0.20% relative decrease.

**FIGURE 33. QUARTERLY NEW INCIDENCE OF SUBSTANCE ABUSE ALONE\* PER 1,000 MEDICARE FFS BENEFICIARIES**




This chart reflects trending of quarterly new incidence of substance abuse alone among Medicare FFS beneficiaries residing in the Toms River community.

\* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.


## Suicide and Intentional Self-Inflicted Injury

**FIGURE 34. SUICIDE AND INTENTIONAL SELF-INFLICTED INJURY PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of suicide and intentional self-inflicted injuries among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 6.49 per 1,000 beneficiaries. After the storm, this rate decreased to 5.96 per 1,000 beneficiaries, which was higher than the county rate. This change reflects an 8.17% relative decrease.

**FIGURE 35. QUARTERLY NEW INCIDENCE OF SUICIDE AND INTENTIONAL SELF-INFLICTED INJURY\* PER 1,000 MEDICARE FFS BENEFICIARIES**



This chart reflects trending of quarterly new incidence of suicide and intentional self-inflicted injuries among Medicare FFS beneficiaries residing in the Toms River community.

\* Quarterly new incidences of conditions that were non-existent (not reported) in the last 12 months.



## RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS

To identify beneficiaries at risk of developing depression or proxy disorders, HQSI conducted a literature review on the potential risk factors for depression or proxy disorders. Previous studies suggested that psychosocial factors, biological factors, deteriorating physical functioning, and medication side effects could increase the risk of depression or proxy disorders.

Based on the literature review and data analysis using factors available through Medicare claims data, the top five risk factors for depression or proxy disorders were identified as: Alzheimer’s disease and related disorders or senile dementia, sleep disturbance, alcohol or substance abuse or tobacco use, hip/pelvic fractures, and amputations (see Appendix B).


For Medicare FFS beneficiaries residing in the community who were diagnosed with these risk factor conditions prior to being diagnosed with depression or proxy disorders, these conditions may have contributed to the risk of developing depression or proxy disorders. The following figures show the prevalence rates for these five conditions in the 12 months before and after Superstorm Sandy.

### Summary

Figure 36. Percent Change of Prevalence of the Top Five Risk Factors of Depression or Proxy Disorders per 1,000 Medicare FFS Beneficiaries						
	Toms River			Ocean		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Any of the Top Five Risk Factors for Depression or Proxy Disorders	170.78	152.85	-10.50	161.01	156.28	-2.94
• Substance or Alcohol Abuse or Tobacco Use	111.70	100.21	-10.29	104.83	102.85	-1.89
• Alzheimer’s Disease and related disorders or Senile Dementia	46.38	35.94	-22.51	39.75	33.91	-14.69
• Sleep Disturbance	24.27	24.35	0.33	25.76	27.77	7.80
• Hip/Pelvic Fractures	8.50	7.09	-16.59	7.72	6.91	-10.49
• Amputations	1.05	0.53	-49.52	1.02	0.74	-27.45

The Toms River community experienced a larger decrease in any of the top five risk factors for depression or proxy disorders than Ocean County. It also experienced a larger decrease in substance or alcohol abuse or tobacco use, Alzheimer’s disease and related disorders or senile demetia, hip/pelvic fractures, and amputations.


**FIGURE 37. ANNUAL PREVALENCE TREND FOR RISK FACTORS OF DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES**



The chart above reflects annual trending in the prevalence of any of the top five risk factors for depression or proxy disorders among Medicare FFS beneficiaries residing in the Toms River community.

## Any of the Top Five Risk Factors for Depression or Proxy Disorders

**FIGURE 38. ANNUAL PREVALENCE OF ANY OF THE TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS PER 1,000 MEDICARE FFS BENEFICIARIES**




The prevalence rate of Medicare FFS beneficiaries residing in the Toms River community with any of the top five risk factors for depression or proxy disorders in the 12 months prior to Superstorm Sandy was 170.78 per 1,000 beneficiaries. After the storm, the rate decreased to 152.85 per 1,000 beneficiaries.

**FIGURE 39. PREVALENCE OF ANY OF THE TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS\* PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES**

October 1, 2011 – September 30, 2012

January 1, 2013 – December 31, 2013




The color-coded map of New Jersey depicts prevalence of any of the top five risk factors from high (red) to low (blue) in the 10 FEMA-declared disaster counties before and after Superstorm Sandy.

\* Mapped using ZIP codes of the 10 counties.

**FIGURE 40. OCEAN COUNTY PREVALENCE OF ANY OF THE TOP FIVE RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS\* PER 1,000 MEDICARE FFS BENEFICIARIES**

October 1, 2011 – September 30, 2012

January 1, 2013 – December 31, 2013




The color-coded map of Ocean County depicts regional variation of prevalence of any of the top five risk factors from high (red) to low (blue) before and after Superstorm Sandy.

\* Mapped using ZIP codes; may not display all the city names located within the ZIP code.

## Alzheimer's Disease and Related Disorders or Senile Dementia


**FIGURE 41. ANNUAL PREVALENCE OF ALZHEIMER'S DISEASE AND RELATED DISORDERS OR SENILE DEMENTIA PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of Medicare FFS beneficiaries residing in the Toms River community with Alzheimer's disease and related disorders or senile dementia in the 12 months prior to Superstorm Sandy was 46.38 per 1,000 beneficiaries. After the storm, the rate decreased to 35.94 per 1,000 beneficiaries, which was higher than the county rate.

## Sleep Disturbance


**FIGURE 42. ANNUAL PREVALENCE OF SLEEP DISTURBANCE PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of Medicare FFS beneficiaries residing in the Toms River community with sleep disturbance in the 12 months prior to Superstorm Sandy was 24.27 per 1,000 beneficiaries. After the storm, the rate increased to 24.35 per 1,000 beneficiaries.

## Substance or Alcohol Abuse or Tobacco Use


**FIGURE 43. ANNUAL PREVALENCE OF SUBSTANCE OR ALCOHOL ABUSE OR TOBACCO USE PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of Medicare FFS beneficiaries residing in the Toms River community with substance or alcohol abuse or tobacco use in the 12 months prior to Superstorm Sandy was 111.70 per 1,000 beneficiaries. After the storm, the rate decreased to 100.21 per 1,000 beneficiaries.

## Hip/Pelvic Fractures


**FIGURE 44. ANNUAL PREVALENCE OF HIP/PELVIC FRACTURES PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of Medicare FFS beneficiaries residing in the Toms River community with hip/pelvic fractures in the 12 months prior to Superstorm Sandy was 8.50 per 1,000 beneficiaries. After the storm, the rate decreased to 7.09 per 1,000 beneficiaries, which was higher than the county rate.

## Amputations

**FIGURE 45. ANNUAL PREVALENCE OF AMPUTATIONS PER 1,000 MEDICARE FFS BENEFICIARIES**



The prevalence rate of Medicare FFS beneficiaries residing in the Toms River community with amputations in the 12 months prior to Superstorm Sandy was 1.05 per 1,000 beneficiaries. After the storm, the rate decreased to 0.53 per 1,000 beneficiaries.

# UTILIZATION OF OUTPATIENT BEHAVIORAL HEALTH SERVICES

## Assessments\*


### Summary

Figure 46. Percent Change of Behavioral Health Service Utilization – Assessments per 1,000 Medicare FFS Beneficiaries						
	Toms River			Ocean		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Annual Depression Screening*	8.78	13.51	53.87	9.50	16.27	71.26
Psychiatric Diagnostic Procedures	57.06	42.43	-25.64	54.39	43.28	-20.43
Neuropsychological Tests	12.31	11.13	-9.59	9.77	12.54	28.35

\* Depression screening comparison time frames are different (January 1, 2012 – December 31, 2012 vs. October 1, 2012 – September 30, 2013)

Annual depression screening in the Toms River community increased from 8.78 per 1,000 beneficiaries before the storm to 13.51 per 1,000 beneficiaries after the storm.

**FIGURE 47. ANNUAL UTILIZATION TREND OF BEHAVIORAL HEALTH ASSESSMENT SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES**



The chart above reflects annual trending in the utilization of behavioral health assessment services among Medicare FFS beneficiaries residing in the Toms River community.

## Depression Screening

One of the long-term goals of this project is to increase the awareness and use of Medicare-covered depression screening among at-risk Medicare FFS beneficiaries residing in the 10 counties during Superstorm Sandy.


\*The utilization rates for Diagnostic Psychological Tests and Health and Behavior Assessment/Intervention are not provided in this profile due to low rates.



Beginning October 2011, depression screening became a Medicare-covered service. According to the CMS Screening for Depression Booklet,<sup>7</sup> Medicare Part B covers an annual screening for depression of 15 minutes in length for beneficiaries in primary care settings when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up. The first quarter of data in this profile for depression screening starts on January 2012 since there were only 14 claims filed for depression screening in the last quarter of 2011 in all 10 counties.


The depression screening utilization rates have been low in all 10 communities.

**FIGURE 48. DEPRESSION SCREENING PER 1,000 MEDICARE FFS BENEFICIARIES**



The rate of depression screening in the Toms River community for calendar year 2012 was 8.78 per 1,000 beneficiaries. After the storm, this rate increased to 13.51 per 1,000 beneficiaries, reflecting a 53.87% relative increase.

**FIGURE 49. QUARTERLY DEPRESSION SCREENING PER 1,000 MEDICARE FFS BENEFICIARIES**



This chart reflects trending of quarterly utilization of depression screening among Medicare FFS beneficiaries residing in the Toms River community.




**Figure 50. Provider Location for Depression Screening\* Claims for Medicare FFS Beneficiaries Residing in Toms River Community**

Providers	1/1/12-12/31/12		1/1/13-12/31/13	
	Number of Claims N=275	Percent	Number of Claims N=449	Percent
Outside of New Jersey	6	2.2	8	1.8
New Jersey	269	97.8	441	98.2
• Ocean County	261	94.9	404	90.0
– Toms River Community	155	56.4	226	50.3
• Other Counties	8	2.9	37	8.2

\* Depression screening is a one time benefit per year. Depression screening comparison time frames are different (January 1, 2012 – December 31, 2012 vs. October 1, 2012 – September 30, 2013)

After Superstorm Sandy, 449 claims for depression screening were filed for beneficiaries residing in the Toms River community, with 1.8% filed outside of New Jersey and 98.2% filed within New Jersey. Of those filed in New Jersey, 90.0% were filed in Ocean County, 50.3% were filed in the Toms River community, and 8.2% were filed in other counties.

**FIGURE 51. DEPRESSION SCREENING\* CLAIMS FOR MEDICARE FFS BENEFICIARIES**




\* Depression screening is a one time benefit per year

During calendar year 2012, 99.27% of depression screening claims recorded in the Toms River community were filed by physicians, 0.36% were filed by nurses, and 0.36% were filed by others. After the storm, 99.55% of depression screening claims were filed by physicians, 0.22% were filed by nurses, and 0.22% were filed by others.

**FIGURE 52. DEPRESSION SCREENING\*  
PER 1,000 MEDICARE FFS BENEFICIARIES IN 10 COUNTIES**

January 1, 2012 – December 31, 2012

January 1, 2013 – December 31, 2013




The color-coded map of New Jersey depicts the use of depression screening from low (red) to high (blue) in the 10 FEMA-declared disaster counties before and after Superstorm Sandy.

\* Mapped using ZIP codes of the 10 counties.

**FIGURE 53. OCEAN COUNTY DEPRESSION SCREENING\*  
PER 1,000 MEDICARE FFS BENEFICIARIES**

January 1, 2012 – December 31, 2012

January 1, 2013 – December 31, 2013




The color-coded map of Ocean County depicts regional variation in the rates of the use of the depression screening benefit from low (red) to high (blue) before and after Superstorm Sandy.

\* Mapped using ZIP codes; may not display all the city names located within the ZIP code.

## Neuropsychological Tests

According to the CMS Mental Health Services Billing Guide, neuropsychological tests are evaluations designed to determine the functional consequences of known or suspected brain injury through testing of the neurocognitive domains responsible for language, perception, memory, learning, problem solving, and adaptation.<sup>8</sup>

**FIGURE 54. NEUROPSYCHOLOGICAL TESTS PER 1,000 MEDICARE FFS BENEFICIARIES**



The rate of neuropsychological tests among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 12.31 per 1,000 beneficiaries. After the storm, this rate decreased to 11.13 per 1,000 beneficiaries, reflecting a 9.59% relative decrease.


**Figure 55. Provider Location for Neuropsychological Tests Claims for Medicare FFS Beneficiaries Residing in Toms River Community\***

Providers	10/1/11-9/30/12		1/1/13-12/31/13	
	Number of Claims N=470	Percent	Number of Claims N=397	Percent
Outside of New Jersey	19	4.0	14	3.5
New Jersey	451	96.0	383	96.5
• Ocean County	372	79.1	295	74.3
– Toms River Community	261	55.5	178	44.8
• Other Counties	79	16.8	88	22.2

\* Number of claims, instead of unique beneficiaries were used in this analysis because a beneficiary can have multiple encounters for these procedures at different locations.

After Superstorm Sandy, 397 claims for neuropsychological tests were filed for beneficiaries residing in the Toms River community, with 3.5% filed outside of New Jersey and 96.5% filed within New Jersey. Of those filed in New Jersey, 74.3% were filed in Ocean County, 44.8% were filed in the Toms River community, and 22.2% were filed in other counties.

**FIGURE 56. NEUROPSYCHOLOGICAL TESTS CLAIMS FOR MEDICARE FFS BENEFICIARIES**




In the 12 months prior to Superstorm Sandy, 73.83% of neuropsychological tests claims were filed by physicians, 24.68% were filed by psychologists, and 1.49% were filed by others. After the storm, 64.48% of neuropsychological tests claims were filed by physicians, 29.47% were filed by psychologists, and 6.05% were filed by others.

### Psychiatric Diagnostic Procedures

According to the CMS Mental Health Services Billing Guide, psychiatric diagnostic evaluation is an integrated biopsychosocial assessment, including history, mental status, and recommendations. The evaluation may include communication with family or other sources and review of diagnostic studies.<sup>8</sup>

**FIGURE 57. PSYCHIATRIC DIAGNOSTIC PROCEDURES PER 1,000 MEDICARE FFS BENEFICIARIES**



The rate of psychiatric diagnostic procedures among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 57.06 per 1,000 beneficiaries. After the storm, this rate decreased to 42.43 per 1,000 beneficiaries. This change reflects a 25.64% relative decrease, a larger decrease than in Ocean County.


**Figure 58. Provider Location for Psychiatric Diagnostic Procedures Claims for Medicare FFS Beneficiaries Residing in Toms River Community\***

Providers	10/1/11-9/30/12		1/1/13-12/31/13	
	Number of Claims N=2,447	Percent	Number of Claims N=1,656	Percent
Outside of New Jersey	74	3.0	57	3.4
New Jersey	2,373	97.0	1,599	96.6
• Ocean County	1,139	46.5	865	52.2
– Toms River Community	758	31.0	607	36.7
• Other Counties	1,234	50.4	734	44.3

\* Number of claims, instead of unique beneficiaries were used in this analysis because a beneficiary can have multiple encounters for these procedures at different locations.

After Superstorm Sandy, 1,656 claims for psychiatric diagnostic procedures were filed for beneficiaries residing in the Toms River community, with 3.4% filed outside of New Jersey and 96.6% filed in New Jersey. Of those filed in New Jersey, 52.2% were filed in Ocean County, 36.7% were filed in the Toms River community, and 44.3% were filed in other counties.

**FIGURE 59. PSYCHIATRIC DIAGNOSTIC PROCEDURES CLAIMS FOR MEDICARE FFS BENEFICIARIES**



In the 12 months prior to Superstorm Sandy, 46.14% of psychiatric diagnostic procedures claims were filed by physicians, 29.42% were filed by psychologists, 12.18% were filed by social workers, 9.40% were filed by nurses, and 2.86% were filed by others.

After the storm, 35.08% of psychiatric diagnostic procedures claims were filed by psychologists, 33.88% were filed by physicians, 13.95% were filed by social workers, 9.60% were filed by nurses, and 7.49% were filed by others.


## Therapies\*

### Summary

Figure 60. Percent Change of Behavioral Health Service Utilization – Therapies per 1,000 Medicare FFS Beneficiaries						
	Toms River			Ocean		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Individual Psychotherapy	46.41	49.05	5.69	48.83	52.21	6.92
Family Psychotherapy	2.32	1.55	-33.19	2.70	2.20	-18.52
Group Psychotherapy	1.28	1.05	-17.97	1.75	1.43	-18.29

The Toms River community experienced a larger decrease in the utilization of family psychotherapy than Ocean County.

**FIGURE 61. ANNUAL UTILIZATION TREND OF BEHAVIORAL HEALTH THERAPY SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES**



This chart reflects trending of annual utilization of behavioral health therapy services among Medicare FFS beneficiaries residing in the Toms River community.


\*The utilization rates for Electroconvulsive Therapy and Biofeedback Therapy are not provided in this profile due to low rates.



## Individual Psychotherapy

According to the CMS Mental Health Services Billing Guide, individual psychotherapy is the treatment of mental illness and behavioral disturbances where the physician or other qualified health professional attempts to alleviate the emotional disturbances, reverse or change maladaptive patterns of behavior, and encourage personality growth and development. This is done through the use of definitive therapeutic communication.<sup>8</sup>

**FIGURE 62. INDIVIDUAL PSYCHOTHERAPY PER 1,000 MEDICARE FFS BENEFICIARIES**




The rate of individual psychotherapy among Medicare FFS beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 46.41 per 1,000 beneficiaries. After the storm, this rate increased to 49.05 per 1,000 beneficiaries, reflecting a 5.69% relative increase.

Figure 63. Provider Location for Individual Psychotherapy Claims for Medicare FFS Beneficiaries Residing in Toms River Community*				
Providers	10/1/11-9/30/12		1/1/13-12/31/13	
	Number of Claims N=9,214	Percent	Number of Claims N=10,327	Percent
Outside of New Jersey	372	4.0	307	3.0
New Jersey	8,842	96.0	10,020	97.0
• Ocean County	5,872	63.7	7,023	68.0
– Toms River Community	3,100	33.6	4,389	42.5
• Other Counties	2,970	32.2	2,997	29.0

\* Number of claims, instead of unique beneficiaries were used in this analysis because a beneficiary can have multiple encounters for these procedures at different locations.

After Superstorm Sandy, 10,327 claims for individual psychotherapy were filed for beneficiaries residing in the Toms River community, with 3.0% filed outside of New Jersey and 97.0% filed in New Jersey. Of those filed in New Jersey, 68.0% were filed in Ocean County, 42.5% were filed in the Toms River community, and 29.0% were filed in other counties.

**FIGURE 64. INDIVIDUAL PSYCHOTHERAPY CLAIMS FOR MEDICARE FFS BENEFICIARIES**




In the 12 months prior to Superstorm Sandy, 46.15% of individual psychotherapy claims were filed by psychologists, 35.45% were filed by social workers, 11.72% were filed by physicians, 4.56% were filed by nurses, and 2.13% were filed by others.

After the storm, 44.24% of individual psychotherapy claims were filed by psychologists, 32.62% were filed by social workers, 14.23% were filed by physicians, 4.74% were filed by nurses, and 4.16% were filed by others.

## Family Psychotherapy

According to the CMS Mental Health Services Billing Guide, family psychotherapy describes the treatment of the family unit when maladaptive behaviors of family members are exacerbating the beneficiary’s mental illness or interfering with treatment. It can also be used to assist the family in addressing the maladaptive behaviors of the patient and improve treatment compliance.<sup>8</sup>

**FIGURE 65. FAMILY PSYCHOTHERAPY PER 1,000 MEDICARE FFS BENEFICIARIES**




The rate of family psychotherapy among beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 2.32 per 1,000 beneficiaries. After the storm, this rate decreased to 1.55 per 1,000 beneficiaries.

Due to these low numbers, no percent change data has been provided for this therapy.

## Group Psychotherapy

According to the CMS Mental Health Services Billing Guide, group psychotherapy is a form of treatment where a selected group of patients are guided by a licensed psychotherapist for the purpose of helping to change maladaptive patterns which interfere with social functioning and are associated with a diagnosable psychiatric illness.<sup>8</sup>

**FIGURE 66. GROUP PSYCHOTHERAPY PER 1,000 MEDICARE FFS BENEFICIARIES**



The rate of group psychotherapy among beneficiaries residing in the Toms River community in the 12 months prior to Superstorm Sandy was 1.28 per 1,000 beneficiaries. After the storm, this rate decreased to 1.05 per 1,000 beneficiaries.

Due to these low numbers, no percent change data has been provided for this therapy.

# INPATIENT HEALTH SERVICES


## Summary

Figure 67. Percent Change of Inpatient Health Service Utilization per 1,000 Medicare FFS Beneficiaries						
	Toms River			Ocean		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Psychiatric Admissions	7.46	7.81	4.69	7.67	6.97	-9.13
Acute Care Hospital Admissions	360.92	318.05	-11.88	334.84	293.55	-12.33
Emergency Department Visits*	281.33	246.63	-12.33	252.79	220.56	-12.75
Observation Stays*	3.91	24.28	520.97	4.48	15.49	245.76

\* Emergency department visits and observation stay rates were based on inpatient Part A claims only.

The Toms River community experienced a larger increase in observation stays than Ocean County.


**FIGURE 68. ANNUAL UTILIZATION TREND OF INPATIENT HEALTH SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES**



This chart reflects trending of annual utilization of inpatient health services among Medicare FFS beneficiaries residing in the Toms River community.

## Psychiatric Hospital Admissions

FIGURE 69. PSYCHIATRIC HOSPITAL ADMISSIONS PER 1,000 MEDICARE FFS BENEFICIARIES




In the 12 months prior to Superstorm Sandy, standalone psychiatric hospitals or distinct part psychiatric units in acute care hospitals in the Toms River community had an admissions rate of 7.46 per 1,000 Medicare FFS beneficiaries. After the storm, the rate increased to 7.81 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 4.69% relative increase.

## Acute Care Hospitals

### Admissions

The following data shows all-cause utilization measures and includes all Medicare FFS beneficiaries, not just beneficiaries with behavioral health conditions.

**FIGURE 70. ACUTE CARE HOSPITAL ADMISSIONS PER 1,000 MEDICARE FFS BENEFICIARIES**




In the 12 months prior to Superstorm Sandy, acute care hospitals in the Toms River community had an acute care admissions rate of 360.92 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 318.05 per 1,000 beneficiaries, which was higher than the county rate. This change reflects an 11.88% relative decrease.

## Observation Stays

According to the U.S. Department of Health and Human Services, observation stays are short-term treatments and assessments provided to Medicare FFS beneficiaries as outpatients to determine whether they require further treatment as inpatients or can be discharged.

**FIGURE 71. OBSERVATION STAYS PER 1,000 MEDICARE FFS BENEFICIARIES**




In the 12 months prior to Superstorm Sandy, observation stays in acute care hospitals in the Toms River community had a rate of 3.91 per 1,000 Medicare FFS beneficiaries. After the storm, the rate increased to 24.28 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 520.97% relative increase, a larger increase than in Ocean County.



## Emergency Department Visits

**FIGURE 72. EMERGENCY DEPARTMENT VISITS PER 1,000 MEDICARE FFS BENEFICIARIES**



In the 12 months prior to Superstorm Sandy, emergency department visits in the Toms River community had a rate of 281.33 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 246.63 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 12.33% relative decrease.

## Within 30 Days of Acute Care Hospital Discharge


### Summary

Figure 73. Percent Change of Inpatient Health Service Utilization Within 30 Days of Discharge per 1,000 Medicare FFS Beneficiaries						
	Toms River			Ocean		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
30-Day Hospital Readmissions	70.24	57.48	-18.17	62.43	50.03	-19.86
Emergency Department Visits*	87.11	73.85	-15.22	78.93	66.11	-16.24
Observation Stays*	9.51	13.31	39.96	7.18	9.82	36.77

\* Emergency department visits and observation stay rates were based on both inpatient and outpatient Part A claims.

The Toms River community experienced a larger increase in observation stays within 30 days of discharge than Ocean County.


**FIGURE 74. ANNUAL UTILIZATION TREND OF INPATIENT HEALTH SERVICES WITHIN 30 DAYS OF DISCHARGE PER 1,000 MEDICARE FFS BENEFICIARIES**



This chart reflects trending of annual utilization of inpatient health services within 30 days of discharge among Medicare FFS beneficiaries residing in the Toms River community.

### 30-Day Hospital Readmissions


**FIGURE 75. 30-DAY HOSPITAL READMISSIONS PER 1,000 MEDICARE FFS BENEFICIARIES**



In the 12 months prior to Superstorm Sandy, acute care hospitals in the Toms River community had a 30-day readmissions rate of 70.24 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 57.48 per 1,000 beneficiaries, which was higher than the county rate. This change reflects an 18.17% relative decrease.

## Observation Stays Within 30 Days of Discharge


**FIGURE 76. OBSERVATION STAYS WITHIN 30 DAYS OF DISCHARGE PER 1,000 MEDICARE FFS BENEFICIARIES**



In the 12 months prior to Superstorm Sandy, the rate of observation stays within 30 days of discharge in the Toms River community was 9.51 per 1,000 Medicare FFS beneficiaries. After the storm, the rate increased to 13.31 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 39.96% relative increase, a larger increase than in Ocean County.

## Emergency Department Visits Within 30 Days of Discharge

**FIGURE 77. EMERGENCY DEPARTMENT VISITS WITHIN 30 DAYS OF DISCHARGE PER 1,000 MEDICARE FFS BENEFICIARIES**



In the 12 months prior to Superstorm Sandy, the rate of emergency department visits within 30 days of discharge in the Toms River community was 87.11 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 73.85 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 15.22% relative decrease.


## Other Settings

### Summary

Figure 78. Percent Change of Other Health Services Utilization per 1,000 Medicare FFS Beneficiaries						
	Toms River			Ocean		
	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change	10/1/11 – 9/30/12	1/1/13 – 12/31/13	% Change
Home Health Agency	112.25	107.95	-3.83	100.09	93.45	-6.63
Skilled Nursing Facility	80.44	70.73	-12.07	76.58	68.46	-10.60
Hospice	32.96	30.73	-6.77	30.49	28.24	-7.38
Medical Rehabilitation	31.56	28.89	-8.46	22.50	20.55	-8.67

The Toms River community experienced a larger decrease in skilled nursing facility services than Ocean County.


**FIGURE 79. ANNUAL UTILIZATION TREND IN OTHER HEALTH SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES**



This chart reflects trending of annual utilization of other health services among Medicare FFS beneficiaries residing in the Toms River community.

## Home Health Agency Services


**FIGURE 80. HOME HEALTH AGENCY SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES**



In the 12 months prior to Superstorm Sandy, the utilization rate of home health agency services in the Toms River community was 112.25 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 107.95 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 3.83% relative decrease.

Skilled Nursing Facility Services

FIGURE 81. SKILLED NURSING FACILITY SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES




In the 12 months prior to Superstorm Sandy, the utilization rate of skilled nursing facility services in the Toms River community was 80.44 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 70.73 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 12.07% relative decrease, a larger decrease than in Ocean County.



## Hospice Services


**FIGURE 82. HOSPICE SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES**



In the 12 months prior to Superstorm Sandy, the utilization rate of hospice services in the Toms River community was 32.96 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 30.73 per 1,000 beneficiaries, which was higher than the county rate. This change reflects a 6.77% relative decrease.

## Medical Rehabilitation Services

**FIGURE 83. MEDICAL REHABILITATION SERVICES PER 1,000 MEDICARE FFS BENEFICIARIES**



In the 12 months prior to Superstorm Sandy, the utilization rate of medical rehabilitation services in the Toms River community was 31.56 per 1,000 Medicare FFS beneficiaries. After the storm, the rate decreased to 28.89 per 1,000 beneficiaries, which was higher than the county rate. This change reflects an 8.46% relative decrease.

## LISTING OF MAJOR HEALTH PROVIDERS

The list below shows the major healthcare facilities that provided 90% of all health services to beneficiaries residing in the Toms River community. These are providers in different care settings and are not restricted to behavioral health providers or services. The map on the following page depicts the location of these providers in relation to the community.

### Acute Care Hospitals

CentraState Medical Center  
 Community Medical Center  
 Deborah Heart and Lung Center  
 Jersey Shore University Medical Center  
 Kimball Medical Center  
 Ocean Medical Center  
 Robert Wood Johnson University Hospital  
 Saint Barnabas Medical Center

### Psychiatric Facilities

Saint Barnabas Behavioral Health Network

### Medical Rehabilitation Centers

HealthSouth Rehabilitation Hospital of Toms River  
 Shore Rehabilitation Institute

### Skilled Nursing Facilities

Arbors Care Center  
 Bey Lea Village Nursing and Rehabilitation Center  
 Community Medical Center Transitional Care Unit  
 Green Acres Manor  
 Hampton Ridge Healthcare and Rehabilitation  
 HealthSouth Rehabilitation Hospital of Toms River  
 Holiday Care Center  
 Manchester Manor Rehabilitation Center  
 Meridian Nursing and Rehabilitation at Brick  
 Meridian Subacute Rehabilitation  
 Rose Garden Nursing and Rehabilitation Center  
 Shore Meadows Rehabilitation and Nursing Center  
 Shorrock Gardens Care Center

### Hospice Facilities


Barnabas Health Van Dyke Hospice & Palliative Care Center at Community Medical Center

### Home Health Agencies

Barnabas Health Home Care – Ocean  
 Holy Redeemer Home Care – New Jersey Shore  
 Meridian Home Care – Ocean County  
 Ocean County Board of Health

## TOMS RIVER COMMUNITY PROVIDERS

The map below shows the major healthcare facilities that served the beneficiaries of the Toms River community based on the Medicare Part A claims database. These are providers in all different care settings and are not restricted to behavioral health providers or services. There are 20 providers located in the community: 10 nursing homes, one hospice, three hospitals, one psychiatric facility, two medical rehabilitation centers and three home health agencies.



## APPENDIX A: BEHAVIORAL HEALTH CONDITIONS

### Documentation and Technical Notes

The following defines the study population, the time frames, and the exclusion and inclusion criteria:

#### *Data Source*

- New Jersey Medicare FFS Part A and Part B claims data and denominator file

#### *Reference Time Period*

- Annual prevalence for the selected behavioral health conditions comparing October 2011 –September 2012 to January 1, 2013 – December 31, 2013
- Annual prevalence trend with quarterly rolling for the selected behavioral health conditions (data starting from January 1, 2011 to December 31, 2013)
- Quarterly new incidence trend of conditions that were not existent (not reported) in the past 12 months of the selected eight behavioral health conditions (data starting from January 1, 2012 to December 31, 2013)

#### *Mapping Tool*

- QGIS Development Team, 2014, QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>
- Source: ZIP code boundaries based on the 2013 U.S. Census Tiger Files

#### *Denominator*

- Denominator was the sum of all eligible Medicare FFS beneficiaries who were in the CMS denominator file during the measurement time frame
- Eligible beneficiaries were computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame
- Where Medicare FFS enrolled days > 0

#### *Numerator*

- Unique Medicare FFS beneficiaries with disease-specific inpatient or outpatient claims during the time frame
- CCW and AHRQ disease diagnosis code match (ICD-9-CM codes) Part A dgns\_cd\_1-25 and dgns\_e\_cd\_1-3; Match Part B dgns\_cd\_1\_12

#### *Exclusions*

- HMO coverage period
- Age <18 or >= 110; Age calculated as end date of time frame or date of death – birth date
- Eligible Medicare FFS days/total measurement days = 0

## Resources

More information on the classification codes, requirements, and processing of the behavioral health conditions highlighted in this profile can be located at the following links:

- Buccaneer, A General Dynamics Company. Chronic Condition Data Warehouse: Additions and Access – Task Order 10 New Clinical Conditions: Requirements and Processing
- [Internet]. [unknown]: Buccaneer, A General Dynamics Company. 2013 May 22 [cited 17 Sep 2013]. Available from: [https://www.ccwdata.org/cs/groups/public/documents/document/clin\\_cond\\_algo\\_req\\_proc.pdf](https://www.ccwdata.org/cs/groups/public/documents/document/clin_cond_algo_req_proc.pdf)
- Healthcare Cost and Utilization Project (H-CUP). Clinical Classifications Software (CCS) for ICD-9-CM [Internet]. Rockville (MD): Agency for Healthcare Research and Quality; Nov 2013 [15 Sep 2013]. Available from: <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>

The following table shows the ICD-9-CM codes for the eight behavioral health conditions:

Behavioral Health Conditions	Numerator: Valid ICD-9-CM Codes
Depression or Proxy Disorders (Depression, Anxiety Disorders or Adjustment Disorders)	29384, 29620, 29621, 29622, 29623, 29624, 29625, 29626, 29630, 29631, 29632, 29633, 29634, 29635, 29636, 30000, 30001, 30002, 30009, 30010, 30020, 30021, 30022, 30023, 30029, 3003, 3004, 3005, 30089, 3009, 3080, 3081, 3082, 3083, 3084, 3089, 3090, 3091, 30922, 30923, 30924, 30928, 30929, 3093, 3094, 30981, 30982, 30983, 30989, 3099, 311, 3130, 3131, 31321, 31322, 3133, 31382, 31383, V790
Depression	29620, 29621, 29622, 29623, 29624, 29625, 29626, 29630, 29631, 29632, 29633, 29634, 29635, 29636, 3004, 311, V790
Anxiety Disorders	29384, 30000, 30001, 30002, 30009, 30010, 30020, 30021, 30022, 30023, 30029, 3003, , 3005, 30089, 3009, 3080, 3081, 3082, 3083, 3084, 3089, 3130, 3131, 31321, 31322, 3133, 31382, 31383
Adjustment Disorders	3090, 3091, 30922, 30923, 30924, 30928, 30929, 3093, 3094, 30981, 30982, 30983, 30989, 3099
Post-Traumatic Stress Disorder (PTSD)	30981
Alcohol or Substance Abuse	2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76072, 76073, 76075, 7795, 96500, 96501, 96502, 96509, V6542 Alcohol Abuse: 2910, 2911, 2912, 2913, 2914, 2915, 2918, 29181, 29182, 29189, 2919, 30300, 30301, 30302, 30303, 30390, 30391, 30392, 30393, 30500, 30501, 30502, 30503, 76071, 9800

Behavioral Health Conditions	Numerator: Valid ICD-9-CM Codes
Substance Abuse Alone	2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76072, 76073, 76075, 7795, 96500, 96501, 96502, 96509, V6542
Suicide and Intentional Self-Inflicted Injury	E9500, E9501, E9502, E9503, E9504, E9505, E9506, E9507, E9508, E9509, E9510, E9511, E9518, E9520, E9521, E9528, E9529, E9530, E9531, E9538, E9539, E954, E9550, E9551, E9552, E9553, E9554, E9555, E9556, E9557, E9559, E956, E9570, E9571, E9572, E9579, E9580, E9581, E9582, E9583, E9584, E9585, E9586, E9587, E9588, E9589, E959, V6284

# APPENDIX B: RISK FACTORS FOR DEPRESSION OR PROXY DISORDERS

## Documentation and Technical Notes

The following defines the study population, the time frame, the exclusion and inclusion criteria, and the literature review references:

### *Data Source*

- New Jersey Medicare FFS Part A and Part B claims data and denominator file

### *Reference Time Period*

- Annual prevalence of risk factors for depression or proxy disorders comparing October 1, 2011 – September 30, 2012 to January 1, 2013 – December 31, 2013
- Annual prevalence trend for risk factors for depression or proxy disorders consists of nine points of data with rolling quarters (starting January 1, 2011 and December 31, 2013)

### *Mapping Tool*

- QGIS Development Team, 2014, QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>
- Source: ZIP code boundaries based on the 2013 U.S. Census Tiger Files

### *Denominator*

- Denominator was the sum of all eligible Medicare FFS beneficiaries who were in the CMS denominator file during the measurement time frame
- Eligible beneficiaries were computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame
- Where Medicare FFS enrolled days > 0

### *Numerator*

- Unique Medicare FFS beneficiaries with disease-specific inpatient or outpatient claims during the time frame
- CCW and AHRQ disease diagnosis code match (ICD-9-CM codes) Part A dgns\_cd\_1-25 and dgns\_e\_cd\_1-3; Match Part B dgns\_cd\_1\_12

### *Exclusions*

- HMO coverage period
- Age <18 or >= 110; Age calculated as end date of time frame or date of death – birth date
- Eligible Medicare FFS days/total measurement days = 0

### *Model*

- Logistic Regression Models were used to determine the top five risk factors with the highest Odds Ratios (OR) ( $p < 0.001$ )



## Resources

More information on the classification codes, requirements, and processing of the combination measure of depression or proxy disorders which includes beneficiaries reported for either depression, anxiety, or adjustment disorders can be located at the following links:

- Buccaneer, A General Dynamics Company. Chronic Condition Data Warehouse: Additions and Access – Task Order 10 New Clinical Conditions: Requirements and Processing
- [Internet]. [unknown]: Buccaneer, A General Dynamics Company. 2013 May 22 [cited 17 Sep 2013]. Available from: [https://www.ccwdata.org/cs/groups/public/documents/document/clin\\_cond\\_algo\\_req\\_proc.pdf](https://www.ccwdata.org/cs/groups/public/documents/document/clin_cond_algo_req_proc.pdf)
- Healthcare Cost and Utilization Project (H-CUP). Clinical Classifications Software (CCS) for ICD-9-CM [Internet]. Rockville (MD): Agency for Healthcare Research and Quality; Nov 2013 [15 Sep 2013]. Available from: <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>

## Literature Review References for Risk Factors for Depression or Proxy Disorders

National Alliance on Mental Illness. Information Helpline: Depression in Older Persons Fact Sheet [Internet]. Arlington (VA): National Alliance on Mental Illness; 2009 Oct [cited 2013 Sep 17]. Available from: <http://www.nami.org/Template.cfm?Section=Helpline1&Template=/ContentManagement/ContentDisplay.cfm&ContentID=144039>

National Institute of Mental Health. Depression: Causes and Risk Factors [Internet]. Bethesda (MD): National Institute of Mental Health; 2013 Jul [cited 2013 Sep 17]. Available from: <http://nihseniorhealth.gov/depression/causesandriskfactors/01.html>

Centers for Disease Control and Prevention and National Association of Chronic Disease Directors. The State of Mental Health and Aging in America [Internet]. Atlanta (GA): National Association of Chronic Disease Directors, 2008 [cited 2013 Sep 19]. 11 p. Available from: [http://www.cdc.gov/aging/pdf/mental\\_health.pdf](http://www.cdc.gov/aging/pdf/mental_health.pdf)

Jacques L, Jensen T, Schafer J, Caplan S, Schott L. Final Coverage Decision Memorandum for Screening for Depression in Adults [Internet]. Baltimore (MD): Centers for Medicare & Medicaid Services; 2011 Oct 14 [cited 2013 Sep 18]. 42 p. Available from: <http://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=251>

Thakur M, Blazer DG. Depression in long-term care. *Journal of the American Medical Directors Association* [Internet]. 2008 Feb [cited 2013 Sep 19];9(2):82-87. Available from: <http://www.amda.com/tools/clinical/depression/DepressioninLongTermCare.pdf>

Sozeri-Varma G. Depression in the elderly: clinical features and risk factors. *Aging and Disease* [Internet]. 2012 Dec [cited 2013 Sep 18];3(6):465-471. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3522513/>

Qian J, Simoni-Wastila L, Rattinger GB, Lehmann S, Langenberg P, et al. Associations of depression diagnosis and antidepressant treatment with mortality among young and disabled Medicare beneficiaries with COPD. *General Hospital Psychiatry*. 2013 Jul 18 [cited 2013 Sep 22]; 35(6):612-618.

## APPENDIX B

---

Shao W, Ahmad R, Khutoryansky N, Aagren M, Bouchard J. Evidence supporting an association between hypoglycemic events and depression. *Current Medical Research and Opinion*. 2013 Sep 23 [cited 2013 Sep 22]: 1-7.

Substance Abuse and Mental Health Services Administration. *The Treatment of Depression in Older Adults: Depression and Older Adults: Key Issues* [Internet]. Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Services Administration, U.S. Department of Health and Human Services, 2011 [cited 2013 Sep 24]. HHS Pub. No. SMA-11-4631. 24 p. Available from: <http://store.samhsa.gov/shin/content/SMA11-4631CD-DVD/SMA11-4631CD-DVD-KeyIssues.pdf>

Himelhoch S, Weller WE, Wu AW, Anderson GF, Cooper LA. Chronic medical illness, depression, and use of acute medical services among Medicare beneficiaries. *Medical Care*. 2004 Jun [cited 2013 Sep 25];42(6):512-521.

Mohile SG, Fan L, Reeve E, Jean-Pierre P, Mustian K, et al. Association of cancer with geriatric syndromes in older Medicare beneficiaries. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* [Internet]. 2011 Apr 10 [cited 2013 Sep 25];29(11): 1458-1464. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3082984/>

Jayadevappa R, Malkowicz SB, Chhatre S, Johnson JC, Gallo JJ. The burden of depression in prostate cancer. *Psycho-oncology*. 2012 Dec [cited 2013 Sep 26];21(12):1338-1345.

Missouri Department of Mental Health. *CPS Facts: Depression and Older Adults* [Internet]. Jefferson City(MO): Missouri Department of Mental Health, [date unknown, cited 2013 Sep 26], 2 p. Available from: <http://dmh.mo.gov/docs/mentalillness/elderlydepress.pdf>

Oregon State University, Washington State University, University of Idaho. *Depression in Later Life: Recognition and Treatment* [Internet]. Corvallis(OR): Pacific Northwest Extension Publication; 2004 Jul [Published April 1990; revised July 2000; cited 2013 Sep 29]; 32 p. Available from: <http://ir.library.oregonstate.edu/xmlui/bitstream/handle/1957/20713/pnw347.pdf>

Cole MG, Dendukuri N. Risk factors for depression among elderly community subjects: a systematic review and meta-analysis. *American Journal of Psychiatry* [Internet]. 2003 Jun [cited 2013 Sep 29]; 160(6):1147-1156. Available from: <http://ajp.psychiatryonline.org/article.aspx?articleid=176272>

Kohn R, Levav I, Garcia ID, Machuca ME, Tamashiro R. Prevalence, risk factors and aging vulnerability for psychopathology following a natural disaster in a developing country. *International Journal of Geriatric Psychiatry*. 2005 Sep [cited 2013 Sep 29];20(9):835-841.

Pietrzak RH, Southwick SM, Tracy M, Galea S, Norris FH. Posttraumatic stress disorder, depression, and perceived needs for psychological care in older persons affected by Hurricane Ike. *Journal of Affective Disorders* [Internet]. 2012 Apr [cited 2013 Sep 30];138(1-2):96-103. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3306486/>

Oriol W. *Psychosocial Issues for Older Adults in Disasters* [Internet]. Washington (DC): Emergency Services and Disaster Relief Branch, Center for Mental Health Services (CMHS), Substance Abuse and Mental Health Services Administration; 1999 [cited 2013 Sep 30]; DHHS Publication No. ESDRB SMA 99-3323. 79 p. Available from: <http://store.samhsa.gov/shin/content/SMA99-3323/SMA99-3323.pdf>

O'Connor EA, Whitlock EP, Gaynes B, Beil TL. Screening for Depression in Adults and Older Adults in Primary Care: An Updated Systematic Review. [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2009 Dec [cited 2013 Sept 30]. 167 p. (Evidence Synthesis No. 75. AHRQ Publication No. 10-05143-EF-1). Available from: <http://www.ncbi.nlm.nih.gov/books/NBK36403/pdf/TOC.pdf>

Noyes K, Liu H, Lyness JM, Friedman B. Medicare beneficiaries with depression: comparing diagnoses in claims data with the results of screening. *Psychiatric Services* [Internet]. 2011 Oct [cited 2013 Sep 30];62(10):1159-1166. Available from: [http://ps.psychiatryonline.org/data/Journals/PSS/4336/pss6210\\_1159.pdf](http://ps.psychiatryonline.org/data/Journals/PSS/4336/pss6210_1159.pdf)

The following table shows the ICD-9-CM codes for the top five risk factors for depression or proxy disorders:

Top Five Risk Factors for Depression or Proxy Disorders*	Numerator: Valid ICD-9-CM Codes
Alzheimer's Disease and Related Disorders or Senile Dementia	3311, 33111, 33119, 3312, 3317, 2900, 29010, 29011, 29012, 29013, 29020, 29021, 2903, 29040, 29041, 29042, 29043, 2940, 2941, 29410, 29411, 2948, 797
Sleep Disturbance	04672, 29182, 29285, 30740, 30741, 30742, 30748, 30749, 32700, 32701, 32702, 32709, 78050, 78051, 78052, 78059
Substance or Alcohol Abuse or Tobacco Use	2910, 2911, 2912, 2913, 2914, 2915, 2918, 29181, 29182, 29189, 2919, 2920, 29211, 29212, 2922, 29281, 29282, 29283, 29284, 29285, 29289, 2929, 30300, 30301, 30302, 30303, 30390, 30391, 30392, 30393, 30400, 30401, 30402, 30403, 30410, 30411, 30412, 30413, 30420, 30421, 30422, 30423, 30430, 30431, 30432, 30433, 30440, 30441, 30442, 30443, 30450, 30451, 30452, 30453, 30460, 30461, 30462, 30463, 30470, 30471, 30472, 30473, 30480, 30481, 30482, 30483, 30490, 30491, 30492, 30493, 30500, 30501, 30502, 30503, 3051, 30510, 30511, 30512, 30513, 30520, 30521, 30522, 30523, 30530, 30531, 30532, 30533, 30540, 30541, 30542, 30543, 30550, 30551, 30552, 30553, 30560, 30561, 30562, 30563, 30570, 30571, 30572, 30573, 30580, 30581, 30582, 30583, 30590, 30591, 30592, 30593, 33392, 3575, 4255, 5353, 53530, 53531, 5710, 5711, 5712, 5713, 64830, 64831, 64832, 64833, 64834, 65550, 65551, 65553, 76071, 76072, 76073, 76075, 7795, 7903, 96500, 96501, 96502, 96509, 9800, V110, V111, V112, V113, V114, V118, V119, V154, V1541, V1542, V1549, V1582, V6285, V6542, V663, V701, V702, V7101, V7102, V7109, V790, V791, V792, V793, V798, V799
Hip/Pelvic Fractures	73314, 73315, 73396, 73397, 73398, 8080, 8081, 8082, 8083, 80841, 80842, 80843, 80849, 80851, 80852, 80853, 80859, 8088, 8089, 82000, 82001, 82002, 82003, 82009, 82010, 82011, 82012, 82013, 82019, 82020, 82021, 82022, 82030, 82031, 82032, 8208, 8209
Amputations	8870, 8871, 8872, 8873, 8874, 8875, 8876, 8877, 8960, 8961, 8962, 8963, 8970, 8971, 8972, 8973, 8974, 8975, 8976, 8977, 9059, 99760, 99761, 99762, 99769

\* Other risk factors for depression or proxy disorders analyzed include Acute Myocardial Infarction (AMI), Stroke/Transient Ischemic Attack, Coronary Artery Bypass Graft Surgery (CABG), Parkinson's Disease, Chronic Obstructive Pulmonary Disease and Bronchiectasis (COPD), Diabetes, Chronic Kidney Disease, Rheumatoid Arthritis/Osteoarthritis (RA/OA), Macular Degeneration, Disability, History of Cancer, Heart Failure, and Acquired Hypothyroidism.

# APPENDIX C: UTILIZATION OF OUTPATIENT MENTAL HEALTH SERVICES

## Documentation and Technical Notes

The following defines the study population, the time frame, and the exclusion and inclusion criteria:

### *Data Source*

- New Jersey Medicare FFS Part A and Part B claims data and denominator file

### *Reference Time Period*

- Annual utilization comparing October 1, 2011 – September 30, 2012 to January 1, 2013 – December 31, 2013
- Annual utilization trend consists of nine points of data with rolling quarters (starting January 1, 2011 and ending December 31, 2013)
- Quarterly utilization trend charts for depression screening contains data from January 1, 2012 to December 31, 2013

### *Mapping Tool*

- QGIS Development Team, 2014, QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>
- Source: ZIP code boundaries based on the 2013 U.S. Census Tiger Files

### *Denominator*

- Denominator was the sum of all eligible Medicare FFS beneficiaries who were in the CMS denominator file during the measurement time frame
- Eligible beneficiaries were computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame
- Where Medicare FFS enrolled days > 0

### *Numerator*

Unique Medicare FFS beneficiaries with specific outpatient mental health service claims

### *Exclusions*

- HMO coverage period
- Age <18 or >= 110; Age calculated as end date of time frame or date of death – birth date
- Eligible Medicare FFS days/total measurement days =0

### *Resources*

More information on the definitions and uses of the outpatient mental health services highlighted in this profile can be located at [http://www.cmsbilling.org/forms/NHIC\\_Medicare\\_B\\_Mental\\_Health\\_billing\\_guide\\_2008.pdf](http://www.cmsbilling.org/forms/NHIC_Medicare_B_Mental_Health_billing_guide_2008.pdf).

The following table shows the CPT/HCPCS codes for the outpatient mental health services:

Mental Health Services	Numerator: CPT/HCPCS Codes
Assessments	
Depression Screening	G0444
Diagnostic Psychological Tests	96101, 96102, 96103, 96105, 96110, 96111
Health and Behavior Assessment/Intervention	96150, 96151, 96152, 96153, 96154, 96155
Neuropsychological Tests	96116, 96118, 96119, 96120
Psychiatric Diagnostic Procedures	90801, 90802, 90791, 90792
Therapies	
Individual Psychotherapy	90804, 90805, 90832, 90833, 90806, 90807, 90834, 90836, 90808, 90809, 90810, 90811, 90812, 90813, 90814, 90815, 90816, 90817, 90818, 90819, 90821, 90822, 90823, 90824, 90826, 90827, 90828, 90829, 90837, 90838, 90839, 90840
Family Psychotherapy	90846, 90847
Group Psychotherapy	90849, 90853, 90857
Electroconvulsive Therapy	90870
Biofeedback Therapy	90901, 90911

# APPENDIX D: UTILIZATION OF SERVICES – INPATIENT AND OTHER SETTINGS

## Documentation and Technical Notes

The following defines the study population, the time frame, and the exclusion and inclusion criteria:

### *Data Source*

New Jersey Medicare FFS Part A claims data and denominator file

### *Reference Time Period*

- Annual utilization comparing October 1, 2011 – September 30, 2012 to January 1, 2013 – December 31, 2013
- Annual utilization trend consists of nine points of data with rolling quarters (starting January 1, 2011 and ending December 31, 2013)

### *Denominator*

- Denominator was the sum of all eligible Medicare FFS beneficiaries who were in the CMS denominator file during the measurement time frame
- Eligible beneficiaries were computed after adjusting for total enrolled FFS days divided by the total measurement days in the time frame
- Where Medicare FFS enrolled days > 0

### *Exclusions*

- HMO coverage period
- Age <18 or >= 110; Age calculated as end date of time frame or date of death – birth date
- Eligible Medicare FFS days/total measurement days =0

### *Utilization Measure*

Refer to Appendix E.

Numerator

Utilization Measure Description	Numerator
Acute Care Hospital Admission	Number of inpatient admissions (Nch_clm_type_cd = 60, 61 )
30-Day Hospital Readmissions	Number of readmissions that occurred within 30 days of hospital discharge (Nch_clm_type_cd = 60, 61)
Emergency Department Visits	Number of emergency department visits, with subsequent inpatient admission (Nch_clm_type_cd = 60, 61 and revenue code in '0450''0451''0452''0456''0459''0981')
Emergency Department Visits within 30 Days of Hospital Discharge	Number of emergency department visits within 30 days of hospital discharge, with or without subsequent admission (Nch_clm_type_cd = 60, 61, 40 and revenue code in '0450''0451''0452''0456' '0459''0981')
Observation Stays	Number of observation stays, with subsequent inpatient admission (Nch_clm_type_cd = 60, 61 and revenue code in '0762')
Observation Stays within 30 Days of Hospital Discharge	Number of observation stays within 30 days of hospital discharge, with or without subsequent admission (Nch_clm_type_cd = 60, 61, 40 and revenue code in '0762')
Home Health Agency Services	Number of eligible beneficiaries with at least one home health agency claim (Nch_clm_type_cd = 10)
Skilled Nursing Facility Services	Number of eligible beneficiaries with at least one skilled nursing facility claim (Nch_clm_type_cd =20, 30)
Hospice Services	Number of eligible beneficiaries with at least one hospice claim (Nch_clm_type_cd = 50)
Medical Rehabilitation Services	Number of eligible beneficiaries with at least one medical rehabilitation claim (Nch_clm_type_cd = 60, 61 and hsp_id format: xxTxxx or between xx3025 and xx3099)
Psychiatric Hospital Admissions	Number of eligible beneficiaries with at least one psychiatric hospital admission claim (Nch_clm_type_cd = 60, 61 and hsp_id format: xxSxxx or between xx4000 and xx4499)

## APPENDIX E: TIME FRAMES AND FORMULAE

Time Frames	
Quarters	Dates
Q1	January 1 to March 31
Q2	April 1 to June 30
Q3	July 1 to September 30
Q4	October 1 to December 31

### Formulae

$$\text{Incidence} = \frac{\text{(Number of unique beneficiaries with new cases during the time frame, condition not present in the past 12 months)}}{\text{(Total unique beneficiaries in the population during the time frame)}}$$

$$\text{Prevalence} = \frac{\text{(Number of unique beneficiaries with the condition during the time frame)}}{\text{(Total unique beneficiaries in the population during the time frame)}}$$

$$\text{Utilization} = \frac{\text{(Number of unique beneficiaries or measures with specific service utilization)}}{\text{(Total unique beneficiaries in the population during the time frame)}}$$

$$\text{Relative change} = \frac{\text{(Current rate-Former rate)}}{\text{(Former rate)}}$$



## APPENDIX F: REFERENCES

1. Weiss MG, Saraceno B, Saxena S, van Ommeren M. Mental health in the aftermath of disasters: consensus and controversy. *The Journal of Nervous and Mental Disease*. 2003 Sep; 191(9):611-615.
2. Foa EB, Stein DJ, McFarlane AC. Symptomatology and psychopathology of mental health problems after disaster. *The Journal of Clinical Psychology* [Internet]. 2006;[cited 16 Sep 2013];67 Suppl 2:15-25.
3. Wang PS, Gruber MJ, Powers RE, Schoenbaum M, Speier AH, Wells KB, Kessler RC. Mental health service use among hurricane Katrina survivors in the eight months after the disaster. *Psychiatry Services* [Internet]. 2007 Nov [cited 16 Sep 2013]; 58(11):1403-1411. Available from: <http://ps.psychiatryonline.org/data/Journals/PSS/3824/07ps1403.pdf>
4. Voelker R. Post-katrina mental health needs prompt group to compile disaster medicine guide. *JAMA*. 2006 Jan [cited 2013 Sep 17]; 295(3):259-260.
5. Centers for Medicare & Medicaid Services. Medicare Claims Database, Parts A and B, January 1, 2011 – March 31, 2013. Baltimore (MD): CMS, Department of Health and Human Services. Accessed: September 15, 2013.
6. U.S. Department of Commerce: United States Census Bureau, American Fact Finder [Internet]. Washington (DC): U.S. Department of Commerce. Median Income in the Past 12 Months (in 2012 Inflation-Adjusted Dollars); 2012 [cited 15 Sep 2013]; [about 2 screens]. Available from: [http://factfinder2.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS\\_12\\_1YR\\_S1903&prodType=table](http://factfinder2.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_12_1YR_S1903&prodType=table)
7. Centers for Medicare & Medicaid Services. Screening for Depression, February 2013. Available from: <https://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-MLN-MLNProducts/Downloads/Screening-for-Depression-Booklet-ICN907799.pdf>
8. Centers for Medicare & Medicaid Services. Mental Health Services Billing Guide, April 2013. Hingham (MA): NHIC, Corp. Apr 2013. 40 p.

## APPENDIX G: PROVIDER SUMMARY TABLES AND PROVIDER LISTINGS

The following defines the data source and time period for the provider summary tables and listings:

### *Data Source*

New Jersey Medicare Part A and Part B FFS claims data

### *Reference Time Period*

Provider summary tables were based on Pre-Sandy time frame (Q1 2011-Q3 2012)

### *Mapping Tool*

ArcGIS Explorer Online. ArcGIS® software by Esri. [www.esri.com](http://www.esri.com)

## **Professional Type by Behavioral Health Services**

The following defines the data source and time period for the provider summary tables and listings:

### *Data Source*

New Jersey Medicare FFS Part B claims data

### *Reference Time Period*

- Professional type of behavioral health service claims during October 1, 2011 – September 30, 2012 and January 1, 2013 – December 31, 2013

### *Professional Type Credentials*

- Physicians: DO, MD
- Psychologists: PhD, PsyD, EdD
- Social Workers: MSW, LCSW
- Nurses: APN, RN, NP
- Others: Other

This Page Left Intentionally Blank

## Executive Summary

Figure 1. Snapshot of Toms River	10
----------------------------------	----

## Demographics

Figure 2. Total Medicare FFS Beneficiaries	11
Figure 3. Percent of Medicare FFS Beneficiaries in the General Population in 2012	11
Figure 4. Percent of Medicare FFS Beneficiary Population by Female	11
Figure 5. Percent of Medicare FFS Beneficiary Population by Race	12
Figure 6. Percent of Medicare FFS Beneficiary Population by Age	12
Figure 7. 2012 Median Household Income (65 years and above)	12

## Prevalence and Incidence

Figure 8. Percent Change of Prevalence of Selected Behavioral Health Conditions	13
Figure 9. Quarterly New Incidence Trend of Selected Behavioral Health Conditions: Depression or Proxy Disorders	14
Figure 10. Quarterly New Incidence Trend of Other Selected Behavioral Health Conditions	14
Figure 11. Annual Prevalence Trend of Selected Behavioral Health Conditions: Depression or Proxy Disorders	15
Figure 12. Annual Prevalence Trend of Other Selected Behavioral Health Conditions	15
Figure 13. Demographics of Depression or Proxy Disorders	16
Figure 14. Depression or Proxy Disorders Rate by Demographic Group	17
Figure 15. Depression or Proxy Disorders Rate by Race	18
Figure 16. Depression or Proxy Disorders Rate by Gender	18
Figure 17. Depression or Proxy Disorders Rate by Age Group	18
Figure 18. Depression or Proxy Disorders (Annual Prevalence, Annual Trend, and Percent Change)	19
Figure 19. Quarterly New Incidence of Depression or Proxy Disorders	19
Figure 20. Prevalence of Depression or Proxy Disorders in 10 Counties	20
Figure 21. Ocean County Prevalence of Depression or Proxy Disorders	21
Figure 22. Depression (Annual Prevalence, Annual Trend, and Percent Change)	22
Figure 23. Quarterly New Incidence of Depression	22
Figure 24. Anxiety Disorders (Annual Prevalence, Annual Trend, and Percent Change)	23
Figure 25. Quarterly New Incidence of Anxiety Disorders	23
Figure 26. Adjustment Disorders (Annual Prevalence, Annual Trend, and Percent Change)	24
Figure 27. Quarterly New Incidence of Adjustment Disorders	24
Figure 28. PTSD (Annual Prevalence, Annual Trend, and Percent Change)	25
Figure 29. Quarterly New Incidence of PTSD	25
Figure 30. Alcohol or Substance Abuse (Annual Prevalence, Annual Trend, and Percent Change)	26
Figure 31. Quarterly New Incidence of Alcohol or Substance Abuse	26
Figure 32. Substance Abuse Alone (Annual Prevalence, Annual Trend, and Percent Change)	27
Figure 33. Quarterly New Incidence of Substance Abuse Alone	27

# INDEX OF FIGURES

---

Figure 34. Suicide and Intentional Self-Inflicted Injury (Annual Prevalence, Annual Trend, and Percent Change)	28
Figure 35. Quarterly New Incidence of Suicide and Intentional Self-Inflicted Injury	28
<b>Risk Factors for Depression or Proxy Disorders</b>	
Figure 36. Percent Change of Prevalence of the Top Five Risk Factors of Depression or Proxy Disorders	29
Figure 37. Annual Prevalence Trend for Risk Factors of Depression or Proxy Disorders	30
Figure 38. Annual Prevalence of Any of the Top Five Risk Factors for Depression or Proxy Disorders	30
Figure 39. Prevalence of Any of the Top Five Risk Factors for Depression or Proxy Disorders in 10 Counties	31
Figure 40. Ocean County Prevalence of Any of the Top Five Risk Factors for Depression or Proxy Disorders	32
Figure 41. Annual Prevalence of Alzheimer’s Disease and Related Disorders or Senile Dementia	33
Figure 42. Annual Prevalence of Sleep Disturbance	33
Figure 43. Annual Prevalence of Substance or Alcohol Abuse or Tobacco Use	33
Figure 44. Annual Prevalence of Hip/Pelvic Fractures	34
Figure 45. Annual Prevalence of Amputations	34
<b>Utilization of Outpatient Behavioral Health Services</b>	
<i>Assessments</i>	
Figure 46. Percent Change of Behavioral Health Service Utilization – Assessments	35
Figure 47. Annual Utilization Trend of Behavioral Health Assessment Services	35
Figure 48. Depression Screening (Annual Utilization, Annual Trend, and Percent Change)	36
Figure 49. Quarterly Depression Screening	36
Figure 50. Provider Location for Depression Screening Claims for Medicare FFS Beneficiaries	37
Figure 51. Depression Screening Claims for Medicare FFS Beneficiaries	37
Figure 52. Depression Screening in 10 Counties	38
Figure 53. Ocean County Depression Screening	39
Figure 54. Neuropsychological Tests (Annual Utilization, Annual Trend, and Percent Change)	40
Figure 55. Provider Location for Neuropsychological Tests Claims for Medicare FFS Beneficiaries	41
Figure 56. Neuropsychological Tests Claims for Medicare FFS Beneficiaries	41
Figure 57. Psychiatric Diagnostic Procedures (Annual Utilization, Annual Trend, and Percent Change)	42
Figure 58. Provider Location for Psychiatric Diagnostic Procedures Claims for Medicare FFS Beneficiaries	43
Figure 59. Psychiatric Diagnostic Procedures Claims for Medicare FFS Beneficiaries	43

## *Therapies*

Figure 60. Percent Change of Behavioral Health Service Utilization – Therapies	44
Figure 61. Annual Utilization Trend of Behavioral Health Therapy Services	44
Figure 62. Individual Psychotherapy (Annual Utilization, Annual Trend, and Percent Change)	45
Figure 63. Provider Location for Individual Psychotherapy Claims for Medicare FFS Beneficiaries	46
Figure 64. Individual Psychotherapy Claims for Medicare FFS Beneficiaries	46
Figure 65. Family Psychotherapy (Annual Utilization and Annual Trend)	47
Figure 66. Group Psychotherapy (Annual Utilization and Annual Trend)	48

## **Inpatient Health Services**

Figure 67. Percent Change of Inpatient Health Service Utilization	49
Figure 68. Annual Utilization Trend of Inpatient Health Services	49
Figure 69. Psychiatric Hospital Admissions (Annual Utilization, Annual Trend, and Percent Change)	50

## *Acute Care Hospitals*

Figure 70. Acute Care Hospital Admissions (Annual Utilization, Annual Trend, and Percent Change)	51
Figure 71. Observation Stays (Annual Utilization, Annual Trend, and Percent Change)	52
Figure 72. Emergency Department Visits (Annual Utilization, Annual Trend, and Percent Change)	53

## *Within 30 Days of Acute Care Hospital Discharge*

Figure 73. Percent Change of Inpatient Health Service Utilization Within 30 Days of Discharge	54
Figure 74. Annual Utilization Trend of Inpatient Health Services Within 30 Days of Discharge	54
Figure 75. 30-Day Hospital Readmissions (Annual Utilization, Annual Trend, and Percent Change)	55
Figure 76. Observation Stays Within 30 Days of Discharge (Annual Utilization, Annual Trend, and Percent Change)	56
Figure 77. Emergency Department Visits Within 30 Days of Discharge (Annual Utilization, Annual Trend, and Percent Change)	57

## *Other Settings*

Figure 78. Percent Change of Other Health Services Utilization	58
Figure 79. Annual Utilization Trend in Other Health Services	58
Figure 80. Home Health Agency Services (Annual Utilization, Annual Trend, and Percent Change)	59
Figure 81. Skilled Nursing Facility Services (Annual Utilization, Annual Trend, and Percent Change)	60
Figure 82. Hospice Services (Annual Utilization, Annual Trend, and Percent Change)	61
Figure 83. Medical Rehabilitation Services (Annual Utilization, Annual Trend, and Percent Change)	62

*This page left intentionally blank*





557 Cranbury Road, Suite 21  
East Brunswick, NJ 08816  
T 732-238-5570 • F 732-238-7766  
[www.hqsi.org](http://www.hqsi.org)